

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11)

**EP 0 702 082 A1**

(12)

**EUROPEAN PATENT APPLICATION**

(43) Date of publication:  
20.03.1996 Bulletin 1996/12

(51) Int. Cl.<sup>6</sup>: **C12N 15/13, C07K 16/42**

(21) Application number: **94115683.8**

(22) Date of filing: **05.10.1994**

(84) Designated Contracting States:  
**AT BE CH DE DK ES FR GB GRIE IT LI LU MC NL  
PT SE**

(30) Priority: **06.10.1993 JP 272950/93**

(71) Applicant: **Hagiwara, Yoshihide  
Takarazuka-shi Hyogo-ken (JP)**

(72) Inventors:  
• **Hagiwara, Hideaki  
Takarazuka-shi, Hyogo-ken (JP)**

• **Aotsuka, Yasuyuki  
Koube-shi, Hyogo-ken (JP)**

(74) Representative: **Weisert, Annekäte, Dipl.-Ing. Dr.-  
Ing. et al  
Patentanwälte  
Kraus Weisert & Partner  
Thomas-Wimmer-Ring 15  
D-80539 München (DE)**

(54) **Amino acid sequences of anti-idiotypic antibodies against anti-cancer human monoclonal antibody, and dna base sequences encoding those sequences**

(57) Amino acid sequences of the H chain and L chain variable regions of mouse monoclonal antibodies Idio 3, Idio 17, Idio 20, Idio 27 and Idio 33 against idiotypes of a cancer cell antigen-specific human immunoglobulin CLN/IgG produced by a human/human fused cell strain CLN/SUZ H11, and base sequences of the genes of the variable regions are disclosed.

The above amino acid sequences and the base sequences are useful in medical and pharmaceutical fields such as prophylaxis, treatment and/or diagnosis of human diseases, and/or in pharmacological and/or biochemical fields, etc. such as biochemical reagents, and reagents for purification of biomacromolecules.

**EP 0 702 082 A1**

## Description

## Detailed Description of the Invention

5 This invention relates to the structure of the variable regions of mouse immunoglobulins against idiotypes of an antigen-specific human immunoglobulin, useful in wide fields, for example in pharmaceutical fields such as prophylaxis, treatment and/or diagnosis of human diseases, and/or in pharmacological and/or biochemical fields such as biochemical reagents and reagents for purification of biomacromolecules.

10 More detailedly, this invention relates to the amino acid sequences of the H chain and L chain variable regions of mouse immunoglobulins against idiotypes of a cancer cell antigen-specific human immunoglobulin produced by a human/human fused cell strain CLN/SUZ H11 from a B cell of a patient carrying human cervical carcinoma and a human lymphoblastoid cell strain, and relates to the base sequences of the genes of the variable regions.

15 Since the development of the technique of formation of monoclonal antibodies by cell fusion or immortalization of cells, many useful antibodies have been obtained using mainly mice. Among them, monoclonal antibodies against malignant tumor cells are utilized not only for fundamental researches such as analyses of tumor antigens, but in serum diagnoses, image diagnoses of tumors using labeled antibodies, and have extremely high utilization value. Particularly, human-derived anti-cancer monoclonal antibodies are expected as ideal antibodies in the clinical field, since they have only faint or no side effects.

20 In such circumstances, one of the present inventors, as disclosed detailedly in Japanese Laid-Open Patent Publication No. 201994/1983 (= U. S. Patent No. 5,286,647; EP-A-839,02157.3), Japanese Laid-Open Patent Publication N. 135898/1984 and Japanese Laid-Open Patent Publication No. 137497/1984, established a cell strain CLN/SUZ H11 (ATCC No. HB 8307) which produces a human monoclonal antibody having a high reactivity with human cancer cells. Interesting findings are obtained about the antibody (named CLN-IgG) produced by this cell strain, that the antibody class is IgG; the isotypes are  $\gamma 1$  type and  $\kappa$  type; and the antibody binds to a cancer antigen immunohistologically existing on the surface of the cancer cells and moreover inhibits proliferation of the cancer cells. At present, the whole amino acid sequence and DNA base sequence of the antibody are clarified (Japanese Laid-Open Patent Publication No. 346792/1992 = WO 92/20799).

30 On the other hand, since Jerne put forward the so-called network theory, various researches have been made on the structure of the variable regions of antibodies. An antibody binds to an antigen at its variable region (antigen combining site). Therefore, the variable regions of antibodies have various three-dimensional-like structures in accordance with the structures of the antigenic determinants on the surfaces of antigens to be recognized. Thus, an antibody itself can be considered to be an antigen, and in the case, the structures of the variable regions of the antibody are called idiotypes, and antibodies against the idiotypes of the antibody are called anti-idiotypic antibodies. The structure corresponding to an antigenic determinant is called an idiotope. An idiotype can be thought to be an aggregate of idiotypes. It was reported that among anti-idiotypic antibodies (Ab2) against an antibody (Ab1) exist antibodies which competitively inhibit binding of Ab1 to an antigen and have idiotopes analogous to antigens recognized by the antibodies, i.e. antibodies having structures as so-called internal images of the antigen.

35 In view of the above findings, anti-idiotypic antibodies are expected to be utilized for the purpose of treatment and/or diagnosis of cancers.

40 For example, as for the purpose of cancer treatment, a vaccine therapy using an anti-idiotypic antibody as an antigen is made possible. It is generally difficult to get cancer antigens in large amounts, and it is restricted from a safety aspect and an ethical aspect to directly immunize human beings with cancer cells as antigens. Therefore, these problems can be avoided by performing immunization with an anti-idiotypic antibody in place of an antigen.

45 In a diagnostic aspect, anti-idiotypic antibodies can be utilized to examine the state of immune reactions against cancer cells. Specifically, it serves for early detection of cancers, judgment of therapeutic effects to detect or determine one's antibodies against cancer antigens existing in the blood or humor of cancer patients.

Under such technical background, problems as stated below are underlying to be solved.

50 1) When anti-idiotypic antibodies are utilized as vaccines or diagnostic drugs, it is necessary to provide these antibodies in large amounts and stably. 2) There is a possibility to give more powerful vaccines or diagnostic drugs abounding in functionality by altering or modifying the antibodies.

A method by gene manipulation is considered as a means for solving the above problems, i.e. a means for realizing improvement of production amount of the antibodies and elevation or modification of the activities of the antibodies.

55 For example, in the case of the problem of 1), it can be considered to solve the problem by cloning such an antibody gene, introducing the gene into host cells such as animal cells or *Escherichia coli*, expressing the antibody gene to give a large amount of the antibody, and in the case of the problem of 2), it can be considered to alter such an antibody so as to have stronger immunogenicity by artificially changing the antibody gene, or to design an antibody molecule having a higher vaccinal activity by adding a function which the antibody does not inherently have, for example an enzymatic activity, an immunity induction activity or the like to the antibody molecule or a fragment thereof.

For accomplishment of these purposes, separation of anti-idiotypic antibody genes, and clarification of their structures are necessary. However, there has not so far been known anything at all about the structures of L chains and H chains constituting anti-idiotypic antibodies against idiotypes of CLN-IgG, and the gene structures of the variable regions having a function to specifically bind to idiotopes of CLN-IgG.

Thus the main object of this invention is to clarify the gene structures of the L chains and the H chains of anti-CLN-IgG idiotype antibodies.

The present inventors have succeeded in creating hybridomas producing, respectively, five kinds of mouse anti-CLN-IgG idiotype antibodies (Idio 3, Idio 17, Idio 20, Idio 27 and Idio 33) having  $\gamma 1$  and  $\kappa$  isotypes against the idiotypes of CLN-IgG; have separated, from the hybridomas, cDNAs encoding the L chains and H chains of the anti-idiotypic antibodies, respectively; have clarified their DNA base sequences; have determined, based on these sequences, the amino acid sequences of the L chains and H chains of the antibodies, respectively; and have completed this invention.

Thus, according to this invention are provided an immunoglobulin H chain variable region fragment which contains a hypervariable region CDR1 having an amino acid sequence selected from

(1) Ser Tyr Trp Met His;  
Asp Tyr Tyr Met Asn; and  
Asn Tyr Trp Met Gln

a hypervariable region CDR2 having an amino acid sequence selected from

(2) Ala Ile Tyr Pro Gly Asn Ser  
Asp Ile Ser Tyr Ser Gln Asn  
Phe Lys Asp;  
Phe Ile Arg Asn Lys Ala  
Asn Leu Tyr Thr Thr Asp  
Tyr Ser Ala Ser Val Lys  
Gly;  
Phe Ile Arg Asn Lys Ala  
Asn Tyr Tyr Thr Thr Glu  
Tyr Ser Ala Ser Val Lys  
Gly; and

Ala Ile Tyr Pro Gly Asp  
Gly Asp Thr Arg Tyr Thr  
Glu Lys Phe Lys Gly

and a hypervariable region CDR3 having an amino acid sequence selected from

- (3) Glu Glu Tyr Asp Tyr Asp  
 Thr Leu Asp Tyr;  
 Asp Arg Gly Gly Arg Asp  
 Trp Tyr Phe Asp Val;  
 Asp Gly Phe Leu Arg Asp  
 Trp Tyr Phe Asp Val; and  
 Ser Gly Tyr Tyr Gly Ser  
 Phe Val Gly Phe Ala Tyr;

and DNA and RNA fragments encoding the immunoglobulin H chain variable region fragment.

According to this invention are further provided an immunoglobulin L chain fragment which contains a hypervariable region CDR1 having an amino acid sequence selected from

- (1) Tyr Arg Ala Ser Lys Ser Val  
 Gln Leu His Leu Ala Ile Val  
 Tyr Met His;  
 Tyr Arg Ala Ser Lys Ser Val  
 Ser Thr Ser Gly Tyr Ser Tyr  
 Met His;  
 Lys Ala Ser Gln Asp Val Asn  
 Thr Ala Val Ala; and  
 Lys Ala Ser Gln Asp Val Thr  
 Thr Asp Val Ala ,

a hypervariable region CDR2 having an amino acid sequence selected from

- (2) Leu Val Ser Asn Leu Glu Ser;  
 Leu Val Ser Asn Leu Asp Ser; and  
 Ser Ala Ser Tyr Arg Tyr Thr,

and a hypervariable region CDR3 having an amino acid sequence selected from

(3) Gln His Ile Arg Val Ala Tyr  
 Thr;  
 Gln His Ile Arg Gly Ala Tyr  
 Thr;  
 Gln His Ile Glu Gly Ala Tyr  
 Thr;  
 Gln Gln His Tyr Ser Pro Pro  
 Leu Thr; and  
 Gln Gln His Tyr Ser Thr Ala  
 Trp Thr;

20 and DNA and RNA fragments encoding the immunoglobulin L chain variable region fragment.

In this invention, cytoplasmic RNAs were prepared from the five mouse hybridomas, respectively; the RNAs were converted to cDNAs by a reverse transcriptase; the antibody genes were amplified using these cDNAs as templates and using the PCR method; the amplified DNA fragments were integrated into plasmids and cloned; the base sequences of the insertion DNAs of the plasmids purified from *Escherichia coli* clones isolated were determined, and the amino acid sequences were determined based on the base sequences. These steps are further detailedly described below.

#### [1] Isolation of cytoplasmic RNAs

Each mouse hybridoma is cultured and proliferated in a culture medium, e.g. and RDF or RPMI 1640 medium, containing 5% fetal bovine serum under a suitable condition, e.g. under a condition of 37°C and a carbon dioxide concentration of 5%; the resultant cells are collected by centrifugation; and the cytoplasmic RNA is extracted from the cells by a conventional method, e.g. a method disclosed in 7.12 of Molecular Cloning (2nd edition, edited by Sambrook et al., Cold Spring Harbor Laboratory Press 1989). The resultant cytoplasmic RNA can further be utilized as a template for cDNA synthesis. Specifically in this invention, the cytoplasmic RNAs were extracted from mouse hybridomas No. 3, No. 17, No. 20, No. 27 and No. 33, and provided for synthesis of cDNAs.

#### [2] Synthesis of cDNAs

Using a cytoplasmic RNA obtained in the step of [1] as a template, a single-strand DNA complementary to the mRNA is synthesized in the presence of dATP, dGTP, dTTP and dCTP using, as a primer, an oligo dT corresponding to a poly A, or a synthetic nucleotide having a random sequence, and a reverse transcriptase. In the specific operations in the invention, cDNAs were synthesized using the cytoplasmic RNAs obtained in the step of [1] as templates and a random hexamer as a primer, respectively, and provided for the step of amplification of the antibody genes.

#### 45 [3] Amplification of antibody genes by PCR

PCR reaction is performed in the presence of dATP, dGTP, dTTP, dCTP and Taq polymerase using as a template a single-strand cDNA obtained in the step of [2] and as a primer a sequence of the antibody gene (e.g., a sequence encoding a constant region, a variable region or a leader region of the antibody gene) to amplify the antibody gene. Suitably in the invention, the antibody genes were amplified using as templates the single-strand cDNAs obtained in the step of [2] and using synthetic DNA oligomers corresponding to the sequences of the leader regions and variable regions of the L chains and H chains of the antibodies, respectively.

#### [4] Cloning of PCR-amplified DNA fragments

55 A PCR-amplified DNA fragment obtained in the step of [3] is, directly or after treatment with restriction enzyme(s), ligated into one of various vectors, for example plasmid vectors such as pUC 18, pCR1000 and pCR™, phage vectors such as M 13 phage, and phagemid vectors such as pUC 118 and pBluescript SK<sup>+</sup> to prepare a vector containing the insertion fragment. Then, *Escherichia coli* is transformed with the vector, and a colony of the *Escherichia coli* containing

the targeted antibody gene fragment is obtained. The purified vector recovered from the *Escherichia coli* is provided as a sample for determination of the DNA base sequence. In the specific operations in the invention, the PCR-amplified DNA fragments obtained in the step of [3] were directly ligated, respectively, into pCR1000 and pCR<sup>TM</sup> plasmid vector; an *Escherichia coli* INVαF<sup>+</sup> was transformed with each of the resultant plasmids; and the plasmids were purified from the resultant *Escherichia coli* colonies, respectively.

#### [5] Determination of the base sequences and amino acid sequences of the DNAs

The base sequence of the DNA at the insertion site in a plasmid obtained in the step of [4] can be determined using the Maxam-Gilbert method or the Sanger method. In the invention, the pCR1000 or pCR<sup>TM</sup> plasmid vectors containing the insertion fragments were purified, respectively; their base sequences were determined by the Sanger method; and the amino acid sequences were presumed based on their base sequences, respectively.

Hereafter, this invention is further specifically described below according to examples.

Drawings referred to in Examples are briefly described as follows.

Fig. 1 is a drawing showing isotypes of monoclonal antibodies Idio 3, Idio 17, Idio 20, Idio 27 and Idio 33.

Fig. 2 is a drawing showing the monoclonal antibodies Idio 3, Idio 17, Idio 20, Idio 27 and Idio 33 specifically bind to CLN-IgG, and do not bind to other human IgGs.

Fig. 3 is a drawing showing that monoclonal antibodies Idio 3, Idio 17, Idio 20, Idio 27 and Idio 33 are competitively inhibiting the binding between CLN-IgG and human matricial carcinoma cell ME-180.

Fig. 4 is a drawing where the amino acid sequences of the H chain variable regions of monoclonal antibodies Idio 3, Idio 17, Idio 20, Idio 27 and Idio 33 are notated in parallel according to the Kabat's notation, and the regions of the hypervariable regions CDR1, CDR2 and CDR3 are determined.

Fig. 5 is a drawing where the amino acid sequences of the L chain variable regions of monoclonal antibodies Idio 3, Idio 17, Idio 20, Idio 27 and Idio 33 are notated in parallel according to the Kabat's notation, and the regions of the hypervariable regions CDR1, CDR2 and CDR3 are determined.

#### Example 1: Preparation of mouse hybridomas

100 µl of 1 mg/ml human IgG (produced by Cappel) is intraperitoneally injected to a Balb/c mouse on the first day after its birth to prepare a mouse having immunological tolerance to human IgG. Six weeks later, the mouse is immunized as follows with CLN-IgG as an antigen.

CLN-IgG purified from a culture medium of a human/human hybridoma CLN/SUZ H11 (ATCC No. HB3307) according to an ammonium sulfate precipitation method and protein A-affinity chromatography was adjusted to a concentration of 2 µg/µl with physiological saline; an equal amount of complete Freund's adjuvant solution was added; and after mixing and emulsification, 100 µl of the emulsion (corresponding to 100 µg of CLN-IgG) was subcutaneously injected into the immunologically tolerated mouse. Thereafter, similar immunization was repeated 4 to 5 times; the murine spleen was enucleated 4 days after the final immunization and made to be spleen cells; and they were used for the following cell fusion.

A mouse parent cells NS-1 (ATCC TIB 18) and the spleen cells are washed with portions of RPMI 1640 medium not containing serum, respectively, and the both of the cells are mixed and centrifuged. 1 ml of 50% polyethylene glycol (average molecular weight: 4,000) is added dropwise to the resultant precipitate over a period of 1 minute. 10 ml of RPMI 1640 medium is further added over a period of 3 minutes, the mixture is centrifuged at 400 x g for 5 minutes, the precipitate is suspended in 10 ml of RPMI 1640 medium containing 20% fetal bovine serum, and the suspension is spread into a 96-well microplate.

Thereafter, the cells were cultured in HAT medium for 14 to 21 days, transferred to HT medium, and finally cultured in RPMI 1640 medium containing 10% fetal bovine serum.

The antibody titers in the culture supernatants on the wells where proliferation was observed were assayed by an enzyme-labeled antibody technique; hybridoma clones secreting monoclonal antibodies which bind to CLN-IgG but not to human IgG were obtained from the appropriate wells by the limiting dilution method; and these hybridoma clones were named No. 3, No. 17, No. 20, No. 27 and No. 33.

#### Example 2: Determination of isotypes of the mouse antibodies

Isotypes of the antibodies secreted from the 5 mouse hybridomas obtained in Example 1 were determined as follows using a mouse monoclonal antibody isotyping kit (produced by Amersham Co.).

The mouse hybridomas are started to be cultured at a concentration each of  $5 \times 10^4$ /ml in portions of RPMI 1640 medium containing 10% fetal bovine serum, respectively, and 5 days later the culture supernatants are obtained, on stick portions of the typing sticks are placed in test tubes, respectively; 3 ml portions of the culture supernatants 5-fold diluted with TBS-T (Tris-buffered saline (TBS, pH 7.6) containing 0.1% Tween 20) are added thereto respectively; and

the mixtures are incubated at room temperature for 15 minutes. The culture supernatants are discarded, 5 ml portions of TBS-T are added, and the typing sticks are washed at room temperature for 5 minutes. TBS-T was discarded, and the washing was repeated once more. 3 ml portions of a peroxidase-labeled anti-mouse antibody 500-fold diluted with TBS-T are added, and the mixtures are incubated at room temperature for 15 minutes. The typing sticks are washed twice in the same manner as above; 3 ml portions of an enzyme substrate solution (obtained by adding one drop of 30% aqueous hydrogen peroxide to 50 ml of a TBS solution of 4-chloro-1-naphtol) are added; the mixtures are subjected to reaction at room temperature for 15 minutes; and then the sticks are washed with distilled water. The isotypes of the mouse antibodies are determined based on the resultant signals, respectively.

As a result, as shown in Fig. 1, all the isotypes of these antibodies were  $\gamma 1$  and  $\kappa$ .

#### Example 3: Examination of specificities of the anti-idiotypic antibodies

It was examined according to a dot blot technique, using an ECL Western blotting detecting reagent (produced by Amersham Co.), that the mouse anti-CLN-IgG idiotype antibodies specifically bind to CLN-IgG. The process is stated below.

CLN-IgG and human IgG1 (produced by Prologen Co.) were diluted with PBS to concentrations of 50 to 0.2  $\mu\text{l/ml}$ , respectively. 2  $\mu\text{l}$  portions of the thus prepared samples were spotted on a number of Hybond-ECL nitrocellulose membrane (produced by Amersham Co.), respectively and after being dried, the nitrocellulose membranes were allowed to stand at room temperature for one hour in PBS-T (0.3% Tween-20-containing PBS) containing 5% skim milk. After being washed with PBS-T, the nitrocellulose membranes were allowed to stand at room temperature for one hour in the culture supernatants (500-fold diluted with PBS-T) of mouse hybridomas No. 3, No. 17, No. 20, No. 27 and No. 33, respectively. After being washed with PBS-T, the nitrocellulose membranes were allowed to stand at room temperature for one hour in portions of a peroxidase-labeled sheep anti-mouse Ig antibody 3,000-fold diluted with PBS-T, respectively. After being washed with PBS-T, the nitrocellulose membranes were subjected to reaction for one minute in portions of the ECL detecting reagent, and sheets of X-ray film were exposed for 30 seconds to the light emitted from the resultant nitrocellulose membranes, respectively.

The results of the sheets of X-ray film developed are shown in Fig. 2. Any of the five antibodies bound to CLN-IgG, but did not bind to human IgG1. Namely, it was revealed that these antibodies are specific to CLN-IgG.

Next, it was examined whether or not the mouse antibodies have an activity to inhibit the binding of a human monoclonal antibody CLN-IgG to a human cancer cell. The method is stated below.

A human cervical carcinoma cell ME-180 (available from ATCC) is cultured in DF medium (a 1:1 mixed-medium of DME:F-12) containing 10% fetal bovine serum. At the stage when the number of the cells becomes  $5 \times 10^5$  to  $1 \times 10^7$ , the cells are detached from the bottom face of the Petri dish using trypsin, collected by centrifugation and sufficiently washed with the medium. A constant number ( $10^5/100 \mu\text{l}$ ) each of the cells is placed in each well of a 96-well microtiter plate, and allowed to stand at  $37^\circ\text{C}$  overnight to be attached on the plate. 50  $\mu\text{l}$  portions of 3% glutaraldehyde solution were added dropwise into the respective wells, and the mixtures are allowed to stand at  $37^\circ\text{C}$  for 20 minutes to fix the cells. The cells of each well are centrifuged at  $200 \times g$  for 10 minutes and washed three times with a gelatin buffer (10 mM phosphate-buffered physiological saline containing 0.3% gelatin); 200  $\mu\text{l}$  portions of 1% bovine serum albumin (BSA) solution are added dropwise, and the mixture is allowed to stand at  $37^\circ\text{C}$  for one hour to block the plate. The cells are washed three times with the gelatin buffer to remove BSA not adsorbed. Thereafter, dilutions at various rates (100 to 1,000,000-fold) of the ascites obtained by intraperitoneally inoculating into mice the various hybridomas secreting the mouse anti-idiotypic antibodies are added dropwise together with CLN-IgG (50  $\mu\text{g}$  each), and the mixtures are subjected to reaction at  $37^\circ\text{C}$  for one hour. The cells of these wells are washed three times with the gelatin buffer. 50  $\mu\text{l}$  portions of a 3,000-fold diluted peroxidase-conjugated goat anti-human Ig antibody (produced by TACO Co.) are added dropwise, respectively, and the mixtures are subjected to reaction at  $37^\circ\text{C}$  for 30 minutes. The cells are washed three times with the gelatin buffer, and portions of a substrate solution containing hydrogen peroxide and o-phenylenediamine are added to perform reaction in a darkroom. 10 minutes later, 50  $\mu\text{l}$  portions of 5N sulfuric acid are added to stop the reaction. When the peroxidase-conjugated goat anti-Ig antibody remains on the microplate, namely when the human IgG to be bound thereto remains, a yellow reaction product having absorption at 490 nm is formed. The amount of CLN-IgG bound to the cancer cell is determined by measuring the amount of the reaction product by a spectrometer.

It was clarified, according to the above method, that all the mouse antibodies Idio 3, Idio 17, Idio 20, Idio 27 and Idio 33 inhibit the binding of CLN-IgG to the cancer cell (Fig. 3).

From the foregoing, these mouse antibodies are antibodies against the idiotypes of CLN-IgG.

#### Example 4: Preparation of RNA

From the five kinds of mouse hybridomas No. 3, No. 17, No. 20, No. 27 and No. 33, the cytoplasmic RNAs were extracted according to the method disclosed in Molecular Cloning (2nd edition, edited by Sambrook et al., Cold Spring Harbor Laboratory Press 1989) 7, 12, as stated below.

10<sup>6</sup> each of the hybridomas cells are collected by centrifugation, and washed twice with 10 times each precipitate's volume of a phosphate-buffered saline. The cells of these groups are centrifuged at 2,000 x g and 4°C for 5 minutes, and the resultant precipitates are suspended in 200 µl portions of an RNA extracting solution (0.14 M NaCl, 1.5 mM MgCl<sub>2</sub>, 10 mM Tris-HCl pH 8.6, 0.5% Nonidet P-40, 1 mM dithiothreitol, 20 mM vanadylribonucleoside complex), respectively. The suspensions are subjected to vortex for 15 seconds and allowed to stand on ice for 5 minutes. The resultant suspensions are centrifuged at 12,000 x g for 30 seconds to remove the cell nuclei as precipitates; to the supernatants are, respectively, added 200 µl portions of a proteinase buffer (0.2 M Tris-HCl pH 8.0, 25 mM EDTA pH 8.0, 0.3 M NaCl, 1.2% SDS) and 1 µl portions of an aqueous proteinase K solution (20 mg/ml); and the mixtures are sufficiently stirred and subjected to incubation at 37°C for 30 minutes. Equal volume portions of phenol/chloroform are added to the reaction solutions, respectively, and the mixtures are stirred, centrifuged at 5,000 x g and room temperature for 10 minutes, and then allowed to separate into organic layers and aqueous layers, respectively. 400 µl portions of isopropanol cooled on ice in advance are added to the aqueous layers recovered, respectively, and the mixtures are allowed to stand on ice for 30 minutes. The mixtures are centrifuged at 12,000 x g and 4°C for 10 minutes to collect RNAs. The resultant RNA precipitates are washed with 1 ml portions of ethanol, dried under reduced pressure and suspended in appropriate amount portions of TE buffer, respectively. Using the cytoplasmic RNAs obtained according to the above operations, the antibody genes are amplified.

#### Example 5: Amplification and cloning of the antibody genes by the RT-PCR method

The antibody genes were amplified from the cytoplasmic RNAs obtained in Example 4, using a GeneAmp® RNA PCR kit (produced by Takara Shuzo Co., Ltd.). First, 20 µl each of reactive solutions were prepared containing PCR buffer II (x1), 5 mM MgCl<sub>2</sub>, 1 mM dATP, 1 mM dGTP, 1 mM dTTP and 1 mM dCTP, 1 U/µl an RNase inhibitor, 2.5 µM a random hexamer, 2.5 U/µl a reverse transcriptase and 100 ng each of the above-mentioned cytoplasmic RNAs, respectively; 20 µl portions of a mineral oil were overlaid thereon respectively; and incubations were performed at room temperature for 10 minutes, at 42°C for 15 minutes, at 99°C for 5 minutes and then at 4°C for 5 minutes to perform cDNA synthesis by reverse transcription reaction. Then, 80 µl portions of a solution consisting of 4 µl of 25 mM MgCl<sub>2</sub>, 8 µl of 10x PCR buffer II, 65.5 µl of sterile distilled water, 0.5 µl of AmpliTaq DNA polymerase (5 U/µl) and 2 µl of PCR primers (each 100 pmoles) were added to the above 20 µl of the reverse transcription reaction solutions; 80 µl portions of the mineral oil were overlaid thereon; and PCR reactions were successfully performed. Each reaction was performed by repeating 30 times the cycle of 94°C for 1.5 minutes, 50°C for 2 minutes and then 72°C for 3 minutes. The base sequences of the PCR primers are shown below. The primers contained in a Ig-Prime™ kit (produced by Novagen Co.) were used except for the primer of the leader sequence C for H chains.

#### Primer for H chains

Leader sequence A	5' GGAATT CATGRASTT SKGGYTMARCTKGRITTT 3'
Leader sequence B	5' GGAATT CATGRAAT GSASCTGGGYWTYCTCTT 3'
Leader sequence C	5' TTAATGGTATCCAGTGT 3'
Constant region	5' CCCAAGCTTCCAGGGRCCARKGGATARACIGRTGG 3'

#### Primer for L chains

Leader sequence A	5' GGAATTGATGRAGWCACAKWCYCAGGTCTTT 3'
Leader sequence B	5' GGAATTGATGGAGACAGACACACTCTCTAT 3'
Constant region	5' CCCAAGCTTACTGGATGGTGGGAAGATGGA 3'

In the above, the alphabets other than A, G, C and T mean the following bases. R=A/G, W=A/T, I=inosine, Y=C/T, D=A/G/T, K=G/T, H=A/C/T, S=C/G, V=A/C/G, M=A/C, B=G/C/T

10 µl portions of the resultant 100 µl each of the PCR reaction products are subjected to 1.5% agarose gel electrophoresis, and it was confirmed that the antibody gene fragments each about 600 bp long were amplified. As a result, in the case of the H chains, the antibody genes derived from No. 3 and No. 17 were amplified in the leader sequence A,



the antibody genes derived from No. 20 and No. 27 were amplified in the leader sequence B, and the antibody gene derived from No. 33 was amplified in the leader sequence C. On the other hand, in the L chains, the antibody genes derived from No. 27 and No. 33 were amplified in the case where the leader sequence A was used, and the antibody genes derived from No. 3, No. 17 and No. 20 were amplified in the leader sequence B.

Each of the PCR-amplified fragments about 600 bp long was integrated into pCR 1000 vector or pCR™ vector using TA cloning kit (produced by Invitrogen Co.). Specifically, ligation mix solutions were prepared by mixing 1 µl portions of the PCR reaction products, 1 µl portions of 10 x the ligation buffer, 2 µl portions of pCR1000 or pCR™ vector (corresponding to 50 µg), 1 µl of T4 DNA ligase and 6 µl portions of sterilized water, respectively; and incubated overnight at 12°C. Separately, 50 µl portions of a suspension of a competent *Escherichia coli* INVαT strain, to which portions were added 2 µl portions of 0.5 M β-mercaptoethanol, respectively, were prepared; and 1 µl portions of the above ligation mix solutions are added thereto, respectively. The mixtures are allowed to stand on ice for 30 minutes, incubated at 42°C for one minute, and rapidly cooled on ice for 2 minutes. 450 µl portions of SOC medium warmed to 42°C in advance were added to the resultant *Escherichia coli* solutions, respectively, and the mixtures are cultured with shaking at 37°C for one hour. Meanwhile, 25 µl portions of X-Gal (40 mg/µl) are spreaded onto a number of LB agar plates each containing Kanamycin (50 µg/ml), respectively, and the agar plates are incubated at 37°C until each X-Gal completely permeates the agar plate.

200 µl portions of the *Escherichia coli* culture broths after completion of culture were spread on the agar plate dried, respectively, and the plates were allowed to stand at 37°C overnight to give white colonies each having Kanamycin resistance.

Plasmids were purified from the *Escherichia coli* clones containing the respective antibody genes, and named 3KB11, 17KB1, 20KB1, 27KA2, 33KA26, 3GB1, 17GB7, 20GA2, 27GA5 and 33GC003, respectively. Purification of the plasmids is performed as follows.

The *Escherichia coli* strains containing the above plasmids, respectively, are cultured 37°C overnight in 100 ml portions of LB medium containing Kanamycin (50 µg/ml), respectively. Each of the resultant culture broths is centrifuged at 3,000 rpm for 10 minutes; the cells collected are suspended in 3 ml of an ice-cooled suspension (50 mM glucose, 10 mM EDTA, 2 mM Tris-HCl pH 8.0); and the suspension is allowed to stand at room temperature for 5 minutes. 6 ml of an alkali lysing solution (0.2 N sodium hydroxide, 1% SDS) is added, and the mixture is mixed by gently turning the centrifugation vessel upside down, and allowed to stand on ice for 5 minutes. 4.5 ml of an ice-cooled neutralizing solution (5 M potassium acetate pH 4.8) is added, and the mixture is centrifuged at 12,000 rpm and 4°C for 10 minutes. The supernatant is transferred into another centrifugation vessel; 1 ml of heat-treated 100 µg/ml RNase A solution is added; and the mixture is subjected to reaction for one hour in an incubator of 37°C to perform RNA digestion. To the reaction solution are added 6 ml of TE buffer-saturated phenol and 6 ml of chloroform/isoamyl alcohol (24:1), and the mixture is subjected to vortex for 30 seconds and then centrifuged at 10,000 rpm and 4°C for 3 minutes. The aqueous layer is transferred into another centrifugation vessel, an equal amount of isopropanol is added, and the mixture is sufficiently mixed and then centrifuged at 10,000 rpm and room temperature for 10 minutes.

The resultant precipitate is washed with 1 ml of 70% cold (-20°C) ethanol, dried under reduced pressure, and dissolved in 480 µl of sterilized water. The solution is transferred into an Eppendorf tube; 120 µl of 4 M NaCl and 600 µl of 13% polyethylene glycol #6000 are added; and the mixture is allowed to stand on ice for 20 minutes. The mixture is then centrifuged at 10,000 rpm and 4°C for 10 minutes, and the precipitate is washed with 1 ml of 70% cold (-20°C) ethanol, dried under reduced pressure and dissolved in 100 µl of TE buffer. The resultant purified plasmid was used as a template for sequencing reaction.

#### Example 6: Determination of the base sequences

Sanger reactions were performed using as templates the plasmids cloning purified in Example 5 and a fluorescence-labeled primer; the reaction products were analyzed by a DNA sequencer DSQ-1 (produced by Shimadzu Corporation); and the DNA base sequences of the insert parts of the plasmids were also determined.

The sequencing reactions were performed using AmpliTaq cycle sequencing kit (produced by Takara Shuzo Co., Ltd.) and a fluorescence-labeled primer in a reagent kit (produced by Wakunaga Pharmaceutical Co., Ltd.) exclusively used for a fluorescence-type DNA sequencer. First, 2 to 4 µg of one of the plasmids purified as stated in Example 5 is mixed with 1 µl of the FITC-labeled primer (1 p mole/µl, forward or reverse is used) and 2 µl of the 10 x cycling mix solution, and sterilized water is added to prepare 10 µl in final volume of a reaction mix. Four tubes are prepared in which 2 µl portions of the termination mix (A, G, C, T) were placed in advance, respectively. 2 µl portions of the above reaction mix were taken and placed into the respective tubes. The mixtures are corrected by centrifugation, 10 µl portions of a mineral oil are overlaid, and cycling reactions are performed under the following conditions: Precycle 95°C, 3 minutes; first cycle 95°C 30 seconds, 60°C 30 seconds, 72°C 1 minute (repeated 15 times); second cycle 95°C 30 seconds, 72°C 1 minute (repeated 15 times); postcycle 4°C.

2 µl portions of a reaction-stopping dye solution (95% formaldehyde, 20 mM EDTA, 0.05% methyl violet) are added, and the mixtures are mixed by centrifugation and preserved at 20°C until they are electrophoresed.

As 5% polyacrylamide gel was used one obtained by adding pur water to 30 g of urea, 6 ml of 10 x TBE buffer (0.89 M Tris-HCl, 0.89 M boric acid, 0.025 M EDTA disodium salt) and 10 ml of 30% acrylamide solution (28.5% acrylamide and 1.5% methylenebisacrylamide, both produced by BIO-RAD Co.) to make the whole volume 60 ml; filtering the mixture with 0.22- $\mu$ m filter; deaerating the filtrate for 30 minutes; adding 150  $\mu$ l of 10% ammonium persulfate and 15  $\mu$ l of TEMEO; allowing the mixture to stand overnight to make it gel.

The gel was set in the DNA sequencer DSQ-1, and prerun was performed at a constant voltage of 1,000 V for one hour. Each of the samples was denatured at 95°C for 3 minutes immediately before electrophoresis, and rapidly cooled on ice, and 2 to 3  $\mu$ l of the reaction solution was sucked up from the bottom part of the tube by a micro-syringe and loaded onto the gel. Samples run was performed at a constant electric power of 20 W for 12 hours.

After completion of electrophoresis, the base sequence was determined using the software attached to DSO-1. The sequence was confirmed by sequencing both of the sense and antisense chains of the same plasmid from both directions.

The resultant base sequences of the variable regions of the H chains and L chains of the five kinds of the mouse monoclonal antibodies, and amino acid sequences presumed therefrom are shown in the following sequence listing. Relation between the sequence numbers and the sequences of the clones are as follows:

Sequence No. 1 : Idio 3 H chain variable region (clone 3GB1)  
 Sequence No. 2 : Idio 17 H chain variable region (clone 17GB7)  
 Sequence No. 3 : Idio 20 H chain variable region (clone 20GA2)  
 Sequence No. 4 : Idio 27 H chain variable region (clone 27GA5)  
 Sequence No. 5 : Idio 33 H chain variable region (clone 33GC003)  
 Sequence No. 6 : Idio 3 L chain variable region (clone 3KB11)  
 Sequence No. 7 : Idio 17 L chain variable region (clone 17KB1)  
 Sequence No. 8 : Idio 20 L chain variable region (clone 20KB1)  
 Sequence No. 9 : Idio 27 L chain variable region (clone 27KA2)  
 Sequence No. 10 : Idio 33 L chain variable region (clone 33KA26)

#### Example 7 Determination of hypervariable regions

The amino acid sequences obtained in Example 6 were notated in parallel according to the numbering of Kabat et al.'s data base (Sequences of proteins of immunological interest Fifth edition, U. S. Department of health and human services, Public health service, National Institutes of Health, NIH Publication No. 91-3242, Kabat et al. 1991), and the amino acid sequences of the hypervariable regions CDR1, CDR2 and CDR3 of each antibody were identified (Fig. 4, H chains, Fig. 5 L chains). In order to confirm the novelty of the identified amino acid sequences of the hypervariable regions CDR1, CDR2 and CDR3, retrieval by a computer was performed using the above Kabat et al.'s data base and a protein data base NBRF-PDB (National Biomedical Research Foundation - protein data base) Release 36.

As a result, the amino acid sequences of Idio 3 H chain CDR1, Idio 17 H chain CDR1, Idio 20 H chain CDR1, Idio 27 H chain CDR1, Idio 33 H chain CDR2, Idio 3 L chain CDR2, Idio 17 L chain CDR2, Idio 27 L chain CDR2 and Idio 33 L chain CDR2 were the same as those of known antibodies, but the amino acid sequences of other CDRs were

revealed to be novel sequences.

## Sequence Listing

Seq. I.D. number : 1

Sequence length : 399

Sequence type : nucleic acid

Strandedness : double

Topology : linear

Sequence kind : mRNA

Original source

Organism : mouse

Sequence characteristics

Symbol expressing characteristics : CDS

Presence position : 1..399

Characteristics determination method : S

Symbol expressing characteristics : sig peptide

Presence position : 1..27

Characteristics determination method : S

## Sequence

CTG TCG GTA ACT TCA GGG GTC TAC TCA GAG GTT CAG CTC CAG CAG TCT 48  
 Leu Ser Val Thr Ser Gly Val Tyr Ser Glu Val Gln Leu Gln Gln Ser

GGG ACT GTG CTG GCA AGG CCT GGG GCT TCA GTG AAG ATG TCC TGC AAG 96  
 Gly Thr Val Leu Ala Arg Pro Gly Ala Ser Val Lys Met Ser Cys Lys

GCT TCG GGC TAC ACC TTT AAC AGC TAG TGG ATG CAC TGG GTA AAA CAG 144  
 Ala Ser Gly Tyr Thr Phe Asn Ser Tyr Trp Met His Trp Val Lys Gln

AGG CCT GGA CAG GGT CTG GAA TGG ATT GGC GCG ATT TAT CCT GGA AAT 192  
 Arg Pro Gly Gln Gly Leu Glu Trp Ile Gly Ala Ile Tyr Pro Gly Asn

AGT GAT ATT AGC TAC AGC CAG AAC TTT AAG GAC AGG GCC AAA CTG ACT 240  
 Ser Asp Ile Ser Tyr Ser Gln Asn Phe Lys Asp Arg Ala Lys Leu Thr

GCC GTC ACA TCC ACC AGC ACT GCC TAC ATG GAA CTC AGA AGC CTG ACA 288  
 Ala Val Thr Ser Thr Ser Thr Ala Tyr Met Glu Leu Arg Ser Leu Thr

AAT GAG GAC TCT GCG GTC TAT TTC TGT ACA AAA GAG GAA TAT GAT TAC 336  
 Asn Glu Asp Ser Ala Val Tyr Phe Cys Thr Lys Glu Glu Tyr Asp Tyr

GAC ACC CTG GAC TAC TGG GGT CAA GGA ACC TCA GTC ACC GTC TCC TCA 384  
 Asp Thr Leu Asp Tyr Trp Gly Gln Gly Thr Ser Val Thr Val Ser Ser

GCC AAA ACG ACA CCC 399  
 Ala Lys Thr Thr Pro

## Sequence Listing

Seq. I.D. number : 2

Sequence length : 402

Sequence type : nucleic acid

Strandedness : double

Topology : linear

Sequence kind : mRNA

Original source

Organism : mouse

## Sequence characteristics

Symbol expressing characteristics : CDS

Presence position : 1..402

Characteristics determination method : S

Symbol expressing characteristics : sig peptide

Presence position : 1..30

Characteristics determination method : S

## Sequence:

```

ATT CTG TCG GTA ACT TCA GGG GTC TAC TCA GAG GTT CAG CTC CAG CAG 48
Ile Leu Ser Val Thr Ser Gly Val Tyr Ser Glu Val Gln Leu Gln Gln
-10          -5          1          5
TCT GGG ACT GTG CTG GCA AGG CCT GGG GCT TCA GTG AAG ATG TCC TGC 96
Ser Gly Thr Val Leu Ala Arg Pro Gly Ala Ser Val Lys Met Ser Cys
10          15          20
AAG GCT TCG GGC TAC ACC TTT AAC AGC TAC TGG ATG CAC TGG GTA AAA 144
Lys Ala Ser Gly Tyr Thr Phe Asn Ser Tyr Trp Met His Trp Val Lys
25          30          35
CAG AGG CCT GGA CAG GGT CTG GAA TGG ATT GGC GCG ATT TAT CCT GGA 192
Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile Gly Ala Ile Tyr Pro Gly
40          45          50
AAT AGT GAT ATT AGC TAC AGC CAG AAC TTT AAG GAC AGG GCC AAA CTG 240
Asn Ser Asp Ile Ser Tyr Ser Gln Asn Phe Lys Asp Arg Ala Lys Leu
55          60          65
ACT GCC GTC ACA TCC ACC AGC ACT GCC TAC ATG GAA CTC AGA AGC CTG 288
Thr Ala Val Thr Ser Thr Ser Thr Ala Tyr Met Glu Leu Arg Ser Leu
70          75          80          85
ACA AAT GAG GAC TCT GCG GTC TAT TTC TGT ACA AAA GAG GAA TAT GAT 336
Thr Asn Glu Asp Ser Ala Val Tyr Phe Cys Thr Lys Glu Glu Tyr Asp
90          95          100
TAC GAC ACC CTG GAC TAC TGG GGT CAA GGA ACC TCA GTC ACC GTC TCC 384
Tyr Asp Thr Leu Asp Tyr Trp Gly Gln Gly Thr Ser Val Thr Val Ser
105          110          115
TCA GCC AAA ACG ACA CCC 402
Ser Ala Lys Thr Thr Pro
120

```

## Sequence Listing

Seq. I.D. number : 3

Sequence length : 438

Sequence type : nucleic acid

Strandedness : double

Topology : linear

Sequence kind : mRNA

Original source

Organism : mouse

## Sequence characteristics

Symbol expressing characteristics : CDS

Presence position : 1..438

Characteristics determination method : S

Symbol expressing characteristics : sig peptide

Presence position : 1..57

Characteristics determination method : S

## Sequence

ATG GAG TTC GGG CTA AAC TGG GTT TTC CTT GTA ACA CTT TTA AAT GGT 248  
Met Glu Phe Gly Leu Asn Trp Val Phe Leu Val Thr Leu Leu Asn Gly 249

-15 -10 -5  
ATC CAG TGT GAG GTG AAG CTG GTG GAG TCT GGA GGA GGC TTG GTA CAG 96  
Ile Gln Cys Glu Val Lys Leu Val Glu Ser Gly Gly Gly Leu Val Gln 97

1 5 10  
CCT GGG GGT TCT CTC AGA CTC TCC TGT GCA ACT TCT GGG TTA ACC TTC 144  
Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Thr Ser Gly Leu Thr Phe 145

15 20 25  
ACT GAT TAC TAC ATG AAC TGG GTC CGC CAG CCT CCA GGA AAG GAA CTT 192  
Thr Asp Tyr Tyr Met Asn Trp Val Arg Gln Pro Pro Gly Lys Glu Leu 193

30 35 40  
GAA TGG TTG GGT TTT ATT AGA AAC AAA GCT AAT CTT TAC ACA ACA GAC 240  
Glu Trp Leu Gly Phe Ile Arg Asn Lys Ala Asn Leu Tyr Thr Thr Asp 241

45 50 55 60  
TAC AGT GCA TCT GTG AAG GGT CGG TTC ACC ATC TCC AGA GAT AAT CCC 288  
Tyr Ser Ala Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Pro 289

65 70 75 80  
CAA AGC ATC CTC TAT CTT CAA ATG AAC ACC CTG ACA ACT GAG GAC AGT 336  
Gln Ser Ile Leu Tyr Leu Gln Met Asn Thr Leu Thr Thr Glu Asp Ser 337

85 90 95 100  
GCC ACT TAT TAG TGT GCA AGA GAT AGG GGG GGG AGG GAC TGG TAC TTC 384  
Ala Thr Tyr Tyr Cys Ala Arg Asp Arg Gly Gly Arg Asp Trp Tyr Phe 385

105 110 115 120  
GAT GTC TGG GGC GCA GGG ACC ACG GTC ACC GTC TCC TCA GCC AAA ACG 432  
Asp Val Trp Gly Ala Gly Thr Thr Val Thr Val Ser Ser Ala Lys Thr 433

125  
ACA CCC  
Thr Pro

125

## Sequence Listing

Seq. I.D. number : 4

Sequence length : 411

Sequence type : nucleic acid

Strandedness : double

Topology : linear

Sequence kind : mRNA

Original source

Organism : mouse

## Sequence characteristics

Symbol expressing characteristics : CDS

Presence position : 1..411

Characteristics determination method : S

Symbol expressing characteristics : signal peptide

Presence position : 1..30

Characteristics determination method : S

## Sequence

```

25  CTT GTA ACA CGT TTA AAT GGT ATC CAG TGT GAG GTG AAG CTG GTG GAG 48
    Leu Val Thr Arg Leu Asn Gly Ile Gln Cys Glu Val Lys Leu Val Glu
    -10          -5          1          5
    TCT GGA GGA GGC TTG GTA CAG CCT GGG GGT TCT CTG AGA CTC TCC TGT 96
    Ser Gly Gly Gly Leu Val Gln Pro Gly Gly Ser Leu Arg Leu Ser Cys
    10          15          20
    GCA ACT TCT GGG TTC ACC TTC ACT GAT TAC TAC ATG AAC TGG GTC CGC 144
    Ala Thr Ser Gly Phe Thr Phe Thr Asp Tyr Tyr Met Asn Trp Val Arg
    25          30          35
    CAG CCT CCA GGA AAG GCA CTT GAG TGG TTG GGT TTT ATT AGA AAC AAA 192
    Gln Pro Pro Gly Lys Ala Leu Glu Trp Leu Gly Phe Ile Arg Asn Lys
    40          45          50
    GCT AAT TAT TAC ACA ACA GAG TAC AGT GCA TCT GTG AAG GGT CGG TTC 240
    Ala Asn Tyr Tyr Thr Thr Glu Tyr Ser Ala Ser Val Lys Gly Arg Phe
    55          60          65
    ACC ATC TCC AGA GAT AAT TCC CAA AGC ATC CTC TAT CTT CAA ATG AAC 288
    Thr Ile Ser Arg Asp Asn Ser Gln Ser Ile Leu Tyr Leu Gln Met Asn
    70          75          80          85
    ACC CTG AGA GGT GAG GAC AGT GCC ACT TAT TAC TGT GCA AGA GAT GGG 336
    Thr Leu Arg Ala Glu Asp Ser Ala Thr Tyr Tyr Cys Ala Arg Asp Gly
    90          95          100
    TTC CTA CGG GAC TGG TAC TTC GAT GTC TGG GGC GCA GGS ACC ACG GTC 384
    Phe Leu Arg Asp Trp Tyr Phe Asp Val Trp Gly Ala Gly Thr Thr Val
    105          110          115
    ACC GTC TCC TCA GCC AAA ACG ACA CCC 411
    Thr Val Ser Ser Ala Lys Thr Thr Pro
    120          125

```

## Sequence Listing

Seq. I.D. number : 5

Sequence length : 363

Sequence type : nucleic acid

Strandedness : double

Topology : linear

Sequence kind : mRNA

Original source

Organism : mouse

## Sequence characteristics

Symbol expressing characteristics : CDS

Presence position : 1..363

Characteristics determination method : S

## Sequence

25	GAG GTT CAG CTC CAG CAG TCT GGG GCT GAA CTG GCA AGA CCT GGG GCT	48
	Glu Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Ala Arg Pro Gly Ala	
	1 5 10 15	
	TCA GTG AAC TTG TCC TGC AAG GCT TCT GGC TAG ACC TTT ACT AAC TAC	96
	Ser Val Asn Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asn Tyr	
30	20 25 30	
	TGG ATG CAG TGG GTA AAA CAG AGG CGT GGA CAG GGT CTG GAA TGG ATT	144
	Trp Met Gln Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile	
	35 40 45	
35	GGG GCT ATT TAT CCT GGA GAT GGT GAT ACT AGG TAC ACT CAG AAG TTC	192
	Gly Ala Ile Tyr Pro Gly Asp Gly Asp Thr Arg Tyr Thr Gln Lys Phe	
	50 55 60	
	AAG GGC AAG GCC ACA TTG ACT GCA GCT AAA TCC TCC AGC ACA GCC TAC	240
40	Lys Gly Lys Ala Thr Leu Thr Ala Ala Lys Ser Ser Ser Thr Ala Tyr	
	65 70 75	
	ATG CAA CTC AGC AGC TTG GCA TCT GAG GAC TCT GCG GTC TAT TAC TGT	288
	Met Gln Leu Ser Ser Leu Ala Ser Glu Asp Ser Ala Val Tyr Tyr Cys	
45	80 85 90 95	
	GCA AGA TCG GGC TAC TAT GGT AGC TTC GTT GGG TTT GCT TAC TGG GGC	336
	Ala Arg Ser Gly Tyr Tyr Gly Ser Phe Val Gly Phe Ala Tyr Trp Gly	
	100 105 110	
50	CAA GGG ACT CTG GTC ACT GTC TCT GCA	363
	Gln Gly Thr Leu Val Thr Val Ser Ala	
	115 120	

## Sequence Listing

Seq. I.D. number : 6

Sequence length : 354

Sequence type : nucleic acid

Strandedness : double

Topology : linear

Sequence kind : mRNA

Original source

Organism : mouse

Sequence characteristics

Symbol expressing characteristics : CDS

Presence position : 1..354

Characteristics determination method : S

## Sequence

GAC ATT GTG CTG ACA CAG TCT CCT GCT TCC TTA GCT GTA TCT CCT CTG	48
Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Pro Leu	
1 5 10 15	
GGG CAG AGG GCC ACC ATC TCA TAC ACG GCC AGC AAA AGT GTG CAG TTA	96
Gly Gln Arg Ala Thr Ile Ser Tyr Arg Ala Ser Lys Ser Val Gln Leu	
20 25 30	
CAT CTG GCT ATA GTT TAT ATG CAC TGG AAC CAA CAG AAA CCA GGA CAG	144
His Leu Ala Ile Val Tyr Met His Trp Asn Gln Gln Lys Pro Gly Gln	
35 40 45	
CCA CCC AGA CTC CTC ATC TAT CTT GTA TCC AAC CTA GAA TCT GGG GTC	192
Pro Pro Arg Leu Leu Ile Tyr Leu Val Ser Asn Leu Glu Ser Gly Val	
50 55 60	
CCT GCC AGG TTC AGT GGC AGT GGG TCT GGG ACA GAC TTC ACC CTC AAC	240
Pro Ala Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Asn	
65 70 75	
ATC CAT CCT GTG GAG GAG GAG GAT GCT CCA ACC TAT TAC TGT CAG CAC	288
Ile His Pro Val Glu Glu Glu Asp Ala Ala Thr Tyr Tyr Cys Gln His	
80 85 90 95	
ATT AGG GTA GCT TAC ACG TTC GGA GGG GGG ACC AAG CTG GAA ATA AAA	336
Ile Arg Val Ala Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys	
100 105 110	
CGG GCT GAT GCT GCA CCA	354
Arg Ala Asp Ala Ala Pro	
115	



## Sequence Listing

Seq. I.D. number : 7

Sequence length : 438

Sequence type : nucleic acid

Strandedness : double

Topology : linear

Sequence kind : mRNA

Original source

Organism : mouse

## Sequence characteristics

Symbol expressing characteristics : CDS

Presence position : 1..438

Characteristics determination method : S

Symbol expressing characteristics : sig peptide

Presence position : 1..39

Characteristics determination method : S

## Sequence

CTA TGG GTA CTG CTG CTC TGG GTT CCA GGT TCC ACT GGT GAC ATT GTG 48  
 Leu Trp Val Leu Leu Trp Val Pro Gly Ser Thr Gly Asp Ile Val  
 -10 -5 1  
 CTG ACA CAG TCT CCT GCT TCC TTA GCT GTA TCT CTG GGG CAG AGG GCC 96  
 Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Leu Gly Gln Arg Ala  
 5 10 15  
 TCC ATC TCA TAC AGG GCC AGC AAA AGT GTC AGT ACA TCT GGC TAT AGT 144  
 Ser Ile Ser Tyr Arg Ala Ser Lys Ser Val Ser Thr Ser Gly Tyr Ser  
 20 25 30  
 TAT ATG CAC TGG AAC CAA CAG AAA CCA GGA CAG CCA CCC AGA CTC CTC 192  
 Tyr Met His Trp Asn Gln Gln Lys Pro Gly Gln Pro Pro Arg Leu Leu  
 35 40 45 50  
 ATC TAT CTT GTA TCC AAC CTA GAA TCT GGG GTC CCT GCC AGG TTC AGT 240  
 Ile Tyr Leu Val Ser Asn Leu Glu Ser Gly Val Pro Ala Arg Phe Ser  
 55 60 65  
 GGC AGT GGG TCT GGG ACA GAC TTC ACC CTC AAC ATC CAT CCT GTG GAG 288  
 Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Asn Ile His Pro Val Glu  
 70 75 80  
 GAG GAG GAT GCT GCA ACC TAT TAC TGT CAG CAC ATT AGG GGA GCT TAC 336  
 Glu Glu Asp Ala Ala Thr Tyr Tyr Cys Gln His Ile Arg Gly Ala Tyr  
 85 90 95  
 ACG TTC GGA GGG GGG ACC AAG CTG GAA ATA AAA CGG GCT GAT GCT GCA 384  
 Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg Ala Asp Ala Ala  
 100 105 110  
 CCA ACT GTA TCC ATC TTC CCA CCA TCC AGT AAG CTT GGG AAA CGG TTC 432  
 Pro Thr Val Ser Ile Phe Pro Pro Ser Ser Lys Leu Gly Lys Arg Phe  
 115 120 125 130  
 GCA CCG 438  
 Ala Pro

## Sequence Listing

Seq. I.D. number : 8  
 Sequence length : 417  
 Sequence type : nucleic acid  
 Strandedness : double  
 Topology : linear  
 Sequence kind : mRNA  
 Original source  
 Organism : mouse

## Sequence characteristics

Symbol expressing characteristics : CDS  
 Presence position : 28..417  
 Characteristics determination method : S  
 Symbol expressing characteristics : sig peptide  
 Presence position : 28..90  
 Characteristics determination method : S

## Sequence

```

25      GGC GCG GTG AGA ACC TTG GGA ATT C ATG GAG ACA GAC ACA CTC CTC      48
          Met Glu Thr Asp Thr Leu Leu
          -20          -15

30      CTA TGG GTA CTG CTG CTC TGG GTT CCA GGT TCC ACT GGT GAC ATT GTG      96
          Leu Trp Val Leu Leu Leu Trp Val Pro Gly Ser Thr Gly Asp Ile Val
          -10          -5          1

          CTG ACA CAG TCT CCT GCT TCC TTA GCA GTT TCT CTC GGG CAG AGG GCC      144
          Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Leu Gly Gln Arg Ala
          5          10          15

35      ACC ATC TCA TAC AGG GCC AGC AAA AGT GTC AGT ACA TCT GGC TAT AGT      192
          Thr Ile Ser Tyr Arg Ala Ser Lys Ser Val Ser Thr Ser Gly Tyr Ser
          20          25          30

          TAT ATG CAC TGG AAC CAA CAG AGA CCA GGA CAG CCA CCC AGA CTC CTC      240
          Tyr Met His Trp Asn Gln Gln Arg Pro Gly Gln Pro Pro Arg Leu Leu
          35          40          45          50

40      ATC TAT CTT GTA TCC AAC CTA GAC TCT GGG GTC CCT GCC AGG TTC AGT      288
          Ile Tyr Leu Val Ser Asn Leu Asp Ser Gly Val Pro Ala Arg Phe Ser
          55          60          65

          GGC AGT GGG TCT GGG ACA GAC TTC ACC CTC AAC ATC CAT CCT GTG GAG      336
          Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Asn Ile His Pro Val Glu
          70          75          80

          GAG GAG GAT GCT GCA ACC TAT TAC TGT CAG CAC ATT GAG GGA GCT TAC      384
          Glu Glu Asp Ala Ala Thr Tyr Tyr Cys Gln His Ile Glu Gly Ala Tyr
          85          90          95

50      ACG TTC GGA GGG GGG ACC AAG CTG GAA ATA AAA      417
          Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
          100          105

```



## Sequence Listing

Seq. I.D. number : 10

Sequence length : 360

Sequence type : nucleic acid

**Strandedness : double**

Topology : linear

Sequence kind : mRNA

**Original source**

Organism : mouse

## Sequence characteristics

Symbol expressing characteristics : CDS

Presence position : 1..360

### Characteristics determination method : S

Symbol expressing characteristics: sig\_peptide

Presence position : 1..12

Characteristics determination method: S

## Sequence

GGT GTT GAC GGA GAC ATT GTG ATG ACA CAG TCT CPC AAA TTC ATG TCC  
Gly Val Asp Gly Asp Ile Val Met Thr Gln Ser His Lys Phe Met Ser

1 5 10

ACA TCA GTT GGA GAC AGG GTC ACC ATC ACC TGC AAG GCC AGT CAG GAT

Thr Ser Val Gly Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp

15 20 25

GTG ACT ACT GAT GTA GCC TGG TAT CAA CAG AAA CCA CGA CAA TCT CCT

Val Thr Thr Asp Val Ala Trp Tyr Gln Gln Lys Pro Arg Gln Ser Pro

30 35 40

AAA CTA CTG ATT TAC TCG GCA TCC TAT CGG TAC ACT GGA GTC CCT GAT

Lys Leu Leu Ile Tyr Ser Ala Ser Tyr Arg Tyr Thr Gly Val Pro Asp

45 50 55

CGC TTC ACT GGC AGT GGA TCT GGG ACG GAT TTC ACT TTC ACC ATC AGC

Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Phe Thr Ile Ser

60 65 70 75

AGT GTG CAG GCT GAA GAC CTG GCA GTT TAT TAC TGT CAG CAA CAT TAT

Ser Val Gln Ala Glu Asp Leu Ala Val Tyr Tyr Cys Gln Gln His Tyr

80 85 90

AGT ACT GCG TGG ACG TTC GGT GGT GGC ACC AAG CTG GAA ATC AAA GCG

Ser Thr Ala Trp Thr Phe Gly Gly Gly Thr Lys Leu Gly Ile Lys Arg

95 100 105

GCT GAT GCT GCA CCA ACT GTA TCC

Ala Asp Ala Ala Pro Thr Val Ser

110 115

110

## SEQUENCE LISTING

5

## (1) GENERAL INFORMATION:

## (i) APPLICANT:

10

- (A) NAME:HAGIWARA, Yoshihide
- (B) STREET:4-14, Hiraisanso
- (C) CITY:Takarazuka-shi
- (D) STATE:Hyogo-ken
- (E) COUNTRY:Japan
- (F) POSTAL CODE (ZIP):none

15

## (ii) TITLE OF INVENTION:AMINO ACID SEQUENCES OF ANTI-IDIOTYPIC ANTIBODIES AGAINST ANTI-CANCER HUMAN MONOCLONAL ANTIBODY, AND DNA BASE SEQUENCES ENCODING THOSE SEQUENCES

20

## (iii) NUMBER OF SEQUENCES:48

## (iv) COMPUTER READABLE FORM:

25

- (A) MEDIUM TYPE:Floppy disk
- (B) COMPUTER:IBM PC compatible
- (C) OPERATING SYSTEM:MS-DOS 4.0
- (D) SOFTWARE:Microsoft Word, Version 5.5

## (v) CURRENT APPLICATION DATA:

- (A) APPLICATION NUMBER:EP - 94 115 683.8
- (B) FILING DATE:October 5, 1994

30

## (2) INFORMATION FOR SEQ ID NO: 1:

## (i) SEQUENCE CHARACTERISTICS:

35

- (A) LENGTH:5 amino acids
- (B) TYPE:amino acid
- (D) TOPOLOGY:linear

## (ii) MOLECULE TYPE:protein

## (ix) FEATURE:

- (A) NAME/KEY:H-CDR1-1
- (D) OTHER INFORMATION:hypervariable region

40

## (xi) SEQUENCE DESCRIPTION:SEQ ID NO:1:

Ser Tyr Trp Met His

5

45

## (2) INFORMATION FOR SEQ ID NO: 2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH:5 amino acids
- (B) TYPE:amino acid
- (D) TOPOLOGY:linear

50

## (ii) MOLECULE TYPE:protein

## (ix) FEATURE:

- (A) NAME/KEY:H-CDR1-2
- (D) OTHER INFORMATION:hypervariable region

## (xi) SEQUENCE DESCRIPTION:SEQ ID NO: 2:

55

Asp Tyr Tyr Met Asn

5

(2) INFORMATION FOR SEQ ID NO: 3:

(1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH:5 amino acids
- (B) TYPE:amino acid
- (D) TOPOLOGY:linear

(ii) MOLECULE TYPE:protein

(ix) FEATURE:

(A) NAME/KEY:H-CDR1-3

(D) OTHER INFORMATION:hypervariable region

(xi) SEQUENCE DESCRIPTION:SEQ ID NO:3:

Asn Tyr Trp Met Gln

5

(2) INFORMATION FOR SEQ ID NO: 4:

(1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH:17 amino acids
- (B) TYPE:amino acid
- (D) TOPOLOGY:linear

(ii) MOLECULE TYPE:protein

(ix) FEATURE:

(A) NAME/KEY:H-CDR2-1

(D) OTHER INFORMATION:hypervariable region

(xi) SEQUENCE DESCRIPTION:SEQ ID NO: 4:

Ala Ile Tyr Pro Gly Asn Ser Asp Ile Ser Tyr Ser Gln Asn Phe Lys

5

10

15

Asp

(2) INFORMATION FOR SEQ ID NO: 5:

(1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH:19 amino acids
- (B) TYPE:amino acid
- (D) TOPOLOGY:linear

(ii) MOLECULE TYPE:protein

(ix) FEATURE:

(A) NAME/KEY:H-CDR2-2

(D) OTHER INFORMATION:hypervariable region

(xi) SEQUENCE DESCRIPTION:SEQ ID NO:5:

Phe Ile Arg Asn Lys Ala Asn Leu Tyr Thr Thr Asp Tyr Ser Ala Ser

5

10

15

Val Lys Gly

(2) INFORMATION FOR SEQ ID NO: 6:

(1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH:19 amino acids
- (B) TYPE:amino acid
- (D) TOPOLOGY:linear

(ii) MOLECULE TYPE:protein

(ix) FEATURE:

(A) NAME/KEY:H-CDR2-3  
(D) OTHER INFORMATION:hypervariable region

(xi) SEQUENCE DESCRIPTION:SEQ ID NO: 6:

Phe Ile Arg Asn Lys Ala Asn Tyr Tyr Thr Thr Glu Tyr Ser Ala Ser  
5 10 15  
Val Lys Gly

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 17 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE:protein

(ix) **FEATURE:**

(A) NAME/KEY: H-CDR2-4

(D) OTHER INFORMATION: hypervariable region

(xi) SEQUENCE DESCRIPTION:SEQ ID NO:7:

Ala Ile Tyr Pro Gly Asp Gly Asp Thr Arg Tyr Thr Glu Lys Phe Lys  
5 10 15  
Gly

(2) INFORMATION FOR SEQ ID NO: 8:

(1) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 10 amino acids

(B) TYPE: amino acid

(D) **TOPOLOGY: linear**

(ii) MOLECULE TYPE:protein

(ix) **FEATURE:**

(A) NAME/KEY: H-CDR3-1

(D) OTHER INFORMATION: hypervariable region

(xi) SEQUENCE DESCRIPTION:SEQ ID NO: 8:

Glu Glu Tyr Asp Tyr Asp Thr Leu Asp Tyr  
5 10

(2) INFORMATION FOR SEQ ID NO: 9:

(1) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 11 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(11) MOLECULE TYPE: protein

(ix) **FEATURE:**

(A) NAME/KEY: H-CDR3-2

(D) OTHER INFORMATION: hypervariable region

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

Asp Arg Gly Gly Arg Asp Trp Tyr Phe Asp Val  
5 10

(2) INFORMATION FOR SEQ ID NO: 10:

(1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH:11 amino acids  
 (B) TYPE:amino acid  
 (D) TOPOLOGY:linear  
 (ii) MOLECULE TYPE:protein  
 (ix) FEATURE:  
 (A) NAME/KEY:H-CDR3-3  
 (D) OTHER INFORMATION:hypervariable region  
 (xi) SEQUENCE DESCRIPTION:SEQ ID NO: 10:

Asp Gly Phe Leu Arg Asp Trp Tyr Phe Asp Val  
 5 10

(2) INFORMATION FOR SEQ ID NO: 11:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH:12 amino acids  
 (B) TYPE:amino acid  
 (D) TOPOLOGY:linear  
 (ii) MOLECULE TYPE:protein  
 (ix) FEATURE:  
 (A) NAME/KEY:H-CDR3-4  
 (D) OTHER INFORMATION:hypervariable region  
 (xi) SEQUENCE DESCRIPTION:SEQ ID NO:11:

Ser Gly Tyr Tyr Gly Ser Phe Val Gly Phe Ala Tyr  
 5 10

(2) INFORMATION FOR SEQ ID NO: 12:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH:17 amino acids  
 (B) TYPE:amino acid  
 (D) TOPOLOGY:linear  
 (ii) MOLECULE TYPE:protein  
 (ix) FEATURE:  
 (A) NAME/KEY:L-CDR1-1  
 (D) OTHER INFORMATION:hypervariable region  
 (xi) SEQUENCE DESCRIPTION:SEQ ID NO: 12:

Tyr Arg Ala Ser Lys Ser Val Gln Leu His Leu Ala Ile Val Tyr Met  
 5 10 15  
 His

(2) INFORMATION FOR SEQ ID NO: 13:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH:16 amino acids  
 (B) TYPE:amino acid  
 (D) TOPOLOGY:linear  
 (ii) MOLECULE TYPE:protein  
 (ix) FEATURE:  
 (A) NAME/KEY:L-CDR1-2  
 (D) OTHER INFORMATION:hypervariable region  
 (xi) SEQUENCE DESCRIPTION:SEQ ID NO:13:

Tyr Arg Ala Ser Lys Ser Val Ser Thr Ser Gly Tyr Ser Tyr Met His  
 5 10 15



## (2) INFORMATION FOR SEQ ID NO: 14:

## (1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH:11 amino acids
- (B) TYPE:amino acid
- (D) TOPOLOGY:linear

(ii) MOLECULE TYPE:protein

## (ix) FEATURE:

- (A) NAME/KEY:L-CDR1-3
- (D) OTHER INFORMATION:hypervariable region

(xi) SEQUENCE DESCRIPTION:SEQ ID NO: 14:

Lys Ala Ser Gln Asp Val Asn Thr Ala Val Ala

5

10

## (2) INFORMATION FOR SEQ ID NO: 15:

## (1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH:11 amino acids
- (B) TYPE:amino acid
- (D) TOPOLOGY:linear

(ii) MOLECULE TYPE:protein

## (ix) FEATURE:

- (A) NAME/KEY:L-CDR1-4
- (D) OTHER INFORMATION:hypervariable region

(xi) SEQUENCE DESCRIPTION:SEQ ID NO:15:

Lys Ala Ser Gln Asp Val Thr Thr Asp Val Ala

5

10

## (2) INFORMATION FOR SEQ ID NO: 16:

## (1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH:7 amino acids
- (B) TYPE:amino acid
- (D) TOPOLOGY:linear

(ii) MOLECULE TYPE:protein

## (ix) FEATURE:

- (A) NAME/KEY:L-CDR2-1
- (D) OTHER INFORMATION:hypervariable region

(xi) SEQUENCE DESCRIPTION:SEQ ID NO: 16:

Leu Val Ser Asn Leu Glu Ser

5

## (2) INFORMATION FOR SEQ ID NO: 17:

## (1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH:7 amino acids
- (B) TYPE:amino acid
- (D) TOPOLOGY:linear

(ii) MOLECULE TYPE:protein

## (ix) FEATURE:

- (A) NAME/KEY:L-CDR2-2
- (D) OTHER INFORMATION:hypervariable region

(xi) SEQUENCE DESCRIPTION:SEQ ID NO:17:

Leu Val Ser Asn Leu Asp Ser

5

(2) INFORMATION FOR SEQ ID NO: 18:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 7 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: protein
- (ix) FEATURE:
- (A) NAME/KEY: L-CDR2-3
  - (D) OTHER INFORMATION: hypervariable region
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:

Ser Ala Ser Tyr Arg Tyr Thr

5

(2) INFORMATION FOR SEQ ID NO: 19:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 8 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: protein
- (ix) FEATURE:
- (A) NAME/KEY: L-CDR3-1
  - (D) OTHER INFORMATION: hypervariable region
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:

Gln His Ile Arg Val Ala Tyr Thr

5

(2) INFORMATION FOR SEQ ID NO: 20:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 8 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: protein
- (ix) FEATURE:
- (A) NAME/KEY: L-CDR3-2
  - (D) OTHER INFORMATION: hypervariable region
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:

Gln His Ile Arg Gly Ala Tyr Thr

5

(2) INFORMATION FOR SEQ ID NO: 21:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 8 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: protein
- (ix) FEATURE:
- (A) NAME/KEY: L-CDR3-3
  - (D) OTHER INFORMATION: hypervariable region

(xi) SEQUENCE DESCRIPTION:SEQ ID NO:21:

Gln His Ile Glu Gly Ala Tyr Thr

5

(2) INFORMATION FOR SEQ ID NO: 22:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH:9 amino acids
- (B) TYPE:amino acid
- (D) TOPOLOGY:linear

(ii) MOLECULE TYPE:protein

(ix) FEATURE:

(A) NAME/KEY:L-CDR3-4

(D) OTHER INFORMATION:hypervariable region

(xi) SEQUENCE DESCRIPTION:SEQ ID NO: 22:

Gln Gln His Tyr Ser Pro Pro Leu Thr

5

(2) INFORMATION FOR SEQ ID NO: 23:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH:9 amino acids
- (B) TYPE:amino acid
- (D) TOPOLOGY:linear

(ii) MOLECULE TYPE:protein

(ix) FEATURE:

(A) NAME/KEY:L-CDR3-5

(D) OTHER INFORMATION:hypervariable region

(xi) SEQUENCE DESCRIPTION:SEQ ID NO:23:

Gln Gln His Tyr Ser Thr Ala Trp Thr

5

(2) INFORMATION FOR SEQ ID NO: 24:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH:34 base pairs
- (B) TYPE:nucleic acid
- (C) STRANDEDNESS:single
- (D) TOPOLOGY:linear

(ii) MOLECULE TYPE:cdna

(iv) ANTISENSE:no

(iii) HYPOTHETICAL:no

(ix) FEATURE:

(A) NAME/KEY:H Leader Sequence A

(D) OTHER INFORMATION:R is A or G;

S is C or G;

K is G or T;

Y is C or T;

M is A or C.

(xi) SEQUENCE DESCRIPTION:SEQ ID NO: 24:

GGGAATTCAT GRASTTSKGG YYTMARCTKG RTTT

34

(2) INFORMATION FOR SEQ ID NO: 25:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH:34 bas pairs  
 (B) TYPE:nucl ic acid  
 (C) STRANDEDNESS:single  
 (D) TOPOLOGY:linear  
 (ii) MOLECULE TYPE:cdna  
 (iii) HYPOTHETICAL:no  
 (iv) ANTISENSE:no  
 (ix) FEATURE:  
 (A) NAME/KEY:H Leader Sequence B  
 (D) OTHER INFORMATION:S is C or G;  
 Y is C or T;  
 W is A or T;  
 R is A or G.  
 (xi) SEQUENCE DESCRIPTION:SEQ ID NO:25:

GGGAATTCAT GRAATGSASC TGGGTYWTYC TCTT

34

## (2) INFORMATION FOR SEQ ID NO: 26:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH:18 base pairs  
 (B) TYPE:nucleic acid  
 (C) STRANDEDNESS:single  
 (D) TOPOLOGY:linear  
 (ii) MOLECULE TYPE:cdna  
 (iii) HYPOTHETICAL:no  
 (iv) ANTISENSE:no  
 (ix) FEATURE:  
 (A) NAME/KEY:H Leader Sequence C  
 (xi) SEQUENCE DESCRIPTION:SEQ ID NO: 26:

TTAAATGGTA TCCAGTGT

18

## (2) INFORMATION FOR SEQ ID NO: 27:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH:35 base pairs  
 (B) TYPE:nucleic acid  
 (C) STRANDEDNESS:single  
 (D) TOPOLOGY:linear  
 (ii) MOLECULE TYPE:cdna  
 (iii) HYPOTHETICAL:no  
 (iv) ANTISENSE:no  
 (ix) FEATURE:  
 (A) NAME/KEY:H Constant Region  
 (D) OTHER INFORMATION:R is A or G;  
 K is G or T;  
 N is inosine.

- (xi) SEQUENCE DESCRIPTION:SEQ ID NO:27:

CCCAAGCTTC CAGGGRCCAR KGGATARACN GRTGG

35

## (2) INFORMATION FOR SEQ ID NO: 28:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH:32 base pairs  
 (B) TYPE:nucleic acid  
 (C) STRANDEDNESS:single  
 (D) TOPOLOGY:linear

- (ii) MOLECULE TYPE:cdna  
 (iii) HYPOTHETICAL:no  
 (iv) ANTISENSE:no  
 (ix) FEATURE:  
     (A) NAME/KEY:L Lead r Sequence A  
     (D) OTHER INFORMATION:R is A or G;  
                                     K is G or T;  
                                     W is A or T;  
                                     Y is C or T.

(xi) SEQUENCE DESCRIPTION:SEQ ID NO: 28:

GGGAATTCAT GRAGWCACAK WCYCAGGTCT TT

32

(2) INFORMATION FOR SEQ ID NO: 29:

- (i) SEQUENCE CHARACTERISTICS:  
     (A) LENGTH:33 base pairs  
     (B) TYPE:nucleic acid  
     (C) STRANDEDNESS:single  
     (D) TOPOLOGY:linear

- (ii) MOLECULE TYPE:cdna  
 (iii) HYPOTHETICAL:no  
 (iv) ANTISENSE:no  
 (ix) FEATURE:  
     (A) NAME/KEY:L Leader Sequence B

(xi) SEQUENCE DESCRIPTION:SEQ ID NO: 29:

GGAATTCAT GGAGACAGAC ACACTCCTGC TAT

33

(2) INFORMATION FOR SEQ ID NO: 30:

- (i) SEQUENCE CHARACTERISTICS:  
     (A) LENGTH:30 base pairs  
     (B) TYPE:nucleic acid  
     (C) STRANDEDNESS:single  
     (D) TOPOLOGY:linear

- (ii) MOLECULE TYPE:cdna  
 (iii) HYPOTHETICAL:no  
 (iv) ANTISENSE:no  
 (ix) FEATURE:  
     (A) NAME/KEY:L constant

(xi) SEQUENCE DESCRIPTION:SEQ ID NO: 30:

CCCAAGCTTA CTGGATGGTG GGAAGATGGA

30

(2) INFORMATION FOR SEQ ID NO: 31:

- (i) SEQUENCE CHARACTERISTICS:  
     (A) LENGTH:357 base pairs  
     (B) TYPE:nucleic acid  
     (C) STRANDEDNESS:double  
     (D) TOPOLOGY:linear

- (ii) MOLECULE TYPE:mRNA  
 (iii) HYPOTHETICAL:no  
 (iv) ANTISENSE:no  
 (vi) ORIGINAL SOURCE:  
     (A) ORGANISM:mouse  
 (ix) FEATURE:

(A) NAME/KEY:Idio 3 H chain variable/Idio 17 H chain variable  
(xi) SEQUENCE DESCRIPTION:SEQ ID NO: 31:

5 GAG GTT CAG CTC GAG CAG TCT GGG ACT GTG CTG GCA AGG CCT GGG GCT 48  
Glu Val Gln Leu Gln Gln Ser Gly Thr Val Leu Ala Arg Pro Gly Ala  
5 10 15

10 TCA GTG AAG ATG TCC TGC AAG GCT TCG GGC TAC ACC TTT AAC AGC TAC 96  
Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Asn Ser Tyr  
20 25 30

15 TGG ATG CAC TGG GTA AAA CAG AGG CCT GGA CAG GGT CTG GAA TGG ATT 144  
Trp Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile  
35 40 45

20 GGC GCG ATT TAT CCT GGA AAT AGT GAT ATT AGC TAC AGC CAG AAC TTT 192  
Gly Ala Ile Tyr Pro Gly Asn Ser Asp Ile Ser Tyr Ser Gln Asn Phe  
50 55 60

25 AAG GAC AGG GCC AAA CTG ACT GCC GTC ACA TCC ACC AGC ACT GCC TAC 240  
Lys Asp Arg Ala Lys Leu Thr Ala Val Thr Ser Thr Ser Thr Ala Tyr  
65 70 75 80

30 ATG GAA CTC AGA AGC CTG ACA AAT GAG GAC TCT GCG GTC TAT TTC TGT 288  
Met Glu Leu Arg Ser Leu Thr Asn Glu Asp Ser Ala Val Tyr Phe Cys  
85 90 95

35 ACA AAA GAG GAA TAT GAT TAC GAC ACC CTG GAC TAC TGG GGT CAA GGA 336  
Thr Lys Glu Glu Tyr Asp Tyr Asp Thr Leu Asp Tyr Trp Gly Gln Gly  
100 105 110

40 ACC TCA GTC ACC GTC TCC TCA 357  
Thr Ser Val Thr Val Ser Ser  
115

(2) INFORMATION FOR SEQ ID NO: 32:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH:366 base pairs.

(B) TYPE:nucleic acid

(C) STRANDEDNESS:double

(D) TOPOLOGY:linear

(ii) MOLECULE TYPE:mRNA

(iii) HYPOTHETICAL:no

(iv) ANTISENSE:no

(vi) ORIGINAL SOURCE:

(A) ORGANISM:mouse

(ix) FEATURE:

(A) NAME/KEY:Idio 20 H chain variable

(xi) SEQUENCE DESCRIPTION:SEQ ID NO: 32:

50 GAG GTG AAG CTG GTG GAG TCT GGA GGA GGC TTG GTA CAG CCT GGG GGT 48  
Glu Val Lys Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly  
5 10 15

55 TCT CTC AGA CTC TCC TGT GCA ACT TCT GGG TTA ACC TTC ACT GAT TAC 96  
Ser Leu Arg Leu Ser Cys Ala Thr Ser Gly Leu Thr Phe Thr Asp Tyr  
20 25 30

5 TAC ATG AAC TGG GTC CGC CAG CCT CCA GGA AAG GAA CTT GAA TGG TTG 144  
Tyr Met Asn Trp Val Arg Gln Pro Pro Gly Lys Glu Leu Glu Trp Leu  
35 40 45

GGT TTT ATT AGA AAC AAA GCT AAT CTT TAC ACA ACA GAC TAC AGT GCA 192  
Gly Phe Ile Arg Asn Lys Ala Asn Leu Tyr Thr Thr Asp Tyr Ser Ala  
50 55 60

10 TCT GTG AAG GGT CGG TTC ACC ATC TCC AGA CAT AAT CCC CAA AGC ATC 240  
Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Pro Gln Ser Ile  
65 70 75 80

CTC TAT CTT CAA ATG AAC ACC CTG ACA ACT GAG GAC AGT GCC ACT TAT 288  
Leu Tyr Leu Gln Met Asn Thr Leu Thr Thr Glu Asp Ser Ala Thr Tyr  
85 90 95

TAC TGT GCA AGA GAT AGG GGG GGG AGG GAC TGG TAC TTC GAT GTC TGG 336  
Tyr Cys Ala Arg Asp Arg Gly Gly Arg Asp Trp Tyr Phe Asp Val Trp  
100 105 110

20 GGC GCA GGG ACC ACG GTC ACC GTC TCC TCA 366  
Gly Ala Gly Thr Thr Val Thr Val Ser Ser  
115 120

## (2) INFORMATION FOR SEQ ID NO: 33:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 366 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: double  
(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: mRNA

## (iii) HYPOTHETICAL: no

## (iv) ANTISENSE: no

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: mouse

## (ix) FEATURE:

- (A) NAME/KEY: Idio 27 H chain variable

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 33:

35 GAG GTG AAG CTG GTG GAG TCT GGA GGA GGC TTG GTA CAG CCT GGG GGT 48  
Glu Val Lys Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly  
5 10 15

40 TCT CTG AGA CTC TCC TGT GCA ACT TCT GGG TTC ACC TTC ACT GAT TAC 96  
Ser Leu Arg Leu Ser Cys Ala Thr Ser Gly Phe Thr Phe Thr Asp Tyr  
20 25 30

45 TAC ATG AAC TGG GTC CGC CAG CCT CCA GGA AAG GCA CTT GAG TGG TTG 144  
Tyr Met Asn Trp Val Arg Gln Pro Pro Gly Lys Ala Leu Glu Trp Leu  
35 40 45

GGT TTT ATT AGA AAC AAA GCT AAT TAT TAC ACA ACA GAG TAC AGT GCA 192  
Gly Phe Ile Arg Asn Lys Ala Asn Tyr Tyr Thr Thr Glu Tyr Ser Ala  
50 55 60

TCT GTG AAG GGT CGG TTC ACC ATC TCC AGA GAT AAT TCC CAA AGC ATC 240  
 S r Val Lys Gly Arg Phe Thr Ile S r Arg Asp Asn S r Gln Ser Ile  
 65 70 75 80

CTC TAT CTT CAA ATG AAC ACC CTG AGA GCT GAG GAC AGT GCC ACT TAT 288  
 Leu Gln Met Asn Thr Leu Thr Leu Arg Ala Glu Asp Ser Ala Thr Tyr  
 85 90 95

TAC TGT GCA AGA GAT GGG TTC CTA CGG GAC TGG TAC TTC GAT GTC TGG 336  
 Tyr Cys Ala Arg Asp Gly Phe Leu Arg Asp Trp Tyr Phe Asp Val Trp  
 100 105 110

GGC GCA GGG ACC ACG GTC ACC GTC TCC TCA 366  
 Gly Ala Gly Thr Thr Val Thr Val Ser Ser  
 115 120

(2) INFORMATION FOR SEQ ID NO: 34:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH:363 base pairs  
 (B) TYPE:nucleic acid  
 (C) STRANDEDNESS:double  
 (D) TOPOLOGY:linear

(ii) MOLECULE TYPE:mRNA

(iii) HYPOTHETICAL:no

(iv) ANTISENSE:no

(vi) ORIGINAL SOURCE:

(A) ORGANISM:mouse

(ix) FEATURE:

(A) NAME/KEY:Idio\_33 H chain variable

(xi) SEQUENCE DESCRIPTION:SEQ ID NO: 34:

GAG GTT CAG CTC CAG CAG TCT GGG GCT GAA CTG GCA AGA CCT GGG GCT 48  
 Glu Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Ala Arg Pro Gly Ala  
 5 10 15

TCA GTG AAC TTG TCC TGC AAG GCT TCT GGC TAC ACC TTT ACT AAC TAC 96  
 Ser Val Asn Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asn Tyr  
 20 25 30

TGG ATG CAG TGG GTA AAA CAG AGG CCT GGA CAG GGT CTG GAA TGG ATT 144  
 Trp Met Gln Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile  
 35 40 45

GGG GCT ATT TAT CCT GGA GAT GGT GAT ACT AGG TAC ACT CAG AAG TTC 192  
 Gly Ala Ile Tyr Pro Gly Asp Gly Asp Thr Arg Tyr Thr Gln Lys Phe  
 50 55 60

AAG GGC AAG GCC ACA TTG ACT GCA GCT AAA TCC TCC AGC ACA GCC TAC 240  
 Lys Gly Lys Ala Thr Leu Thr Ala Ala Lys Ser Ser Ser Thr Ala Tyr  
 65 70 75 80

ATG CAA CTC ACC AGC TTG GCA TCT GAG GAC TCT GCG GTC TAT TAC TGT 288  
 Met Gln Leu Ser Ser Leu Ala Ser Glu Asp Ser Ala Val Tyr Tyr Cys  
 85 90 95



GCA AGA TCG GGC TAC TAT GGT AGC TTC GTT GGG TTT GCT TAC TGG GGC 336  
Ala Arg Ser Gly Tyr Tyr Gly S r Phe Val Gly Phe Ala Tyr Trp Gly  
100 105 110

CAA GGG ACT CTG GTC ACT GTC TCT GCA 363  
Gln Gly Thr Leu Val Thr Val Ser Ala  
115 120

(2) INFORMATION FOR SEQ ID NO: 35:

(1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH:336 base pairs
- (B) TYPE:nucleic acid
- (C) STRANDEDNESS:double
- (D) TOPOLOGY:linear

(ii) MOLECULE TYPE:mRNA

(iii) HYPOTHETICAL:no

(iv) ANTISENSE:no

(vi) ORIGINAL SOURCE:

(A) ORGANISM:mouse

(ix) FEATURE:

(A) NAME/KEY:Idio 3 L chain variable

(xi) SEQUENCE DESCRIPTION:SEQ ID NO: 35:

GAC ATT GTG CTG ACA CAG TCT CCT GCT TCC TTA GCT GTA TCT CCT CTG 48  
Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Pro Leu  
5 10 15

GGG CAG AGG GCC ACC ATC TCA TAC AGG GCC AGC AAA AGT GTG CAG TTA 96  
Gly Gln Arg Ala Thr Ile Ser Tyr Arg Ala Ser Lys Ser Val Gln Leu  
20 25 30

CAT CTG GCT ATA GTT TAT ATG CAC TGG AAC CAA CAG AAA CCA GGA CAG 144  
His Leu Ala Ile Val Tyr Met His Trp Asn Gln Gln Lys Pro Gly Gln  
35 40 45

CCA CCC AGA CTC CTC ATC TAT CTT GTA TCC AAC CTA GAA TCT GGG GTC 192  
Pro Pro Arg Leu Leu Ile Tyr Leu Val Ser Asn Leu Glu Ser Gly Val  
50 55 60

CCT GCC AGG TTC AGT GGC AGT GGG TCT GGG ACA GAC TTC ACC CTC AAC 240  
Pro Ala Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Asn  
65 70 75 80

ATC CAT CCT GTG GAG GAG GAG GAT GCT GCA ACC TAT TAC TGT CAG CAC 288  
Ile His Pro Val Glu Glu Glu Asp Ala Ala Thr Tyr Tyr Cys Gln His  
85 90 95

ATT AGG GTA GCT TAC ACG TTC GGA GGG GGG ACC AAG CTG GAA ATA AAA 336  
Ile Arg Val Ala Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys  
100 105 110

(2) INFORMATION FOR SEQ ID NO: 36:

(1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH:330 base pairs

(B) TYPE:nucleic acid  
 (C) STRANDEDNESS:double  
 (D) TOPOLOGY:linear  
 (ii) MOLECULE TYPE:mRNA  
 (iii) HYPOTHETICAL:no  
 (iv) ANTISENSE:no  
 (vi) ORIGINAL SOURCE:  
 (A) ORGANISM:mouse  
 (ix) FEATURE:  
 (A) NAME/KEY:Idio 17 L chain variable  
 (xi) SEQUENCE DESCRIPTION:SEQ ID NO: 36:

GAC ATT GTG CTG ACA CAG TCT CCT GCT TCC TTA GCT GTA TCT CTG GGG 48  
 Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Leu Gly  
 5 10 15  
 CAG AGG GCC TCC ATC TCA TAC AGG GCC AGC AAA AGT GTC AGT ACA TCT 96  
 Gln Arg Ala Ser Ile Ser Tyr Arg Ala Ser Lys Ser Val Ser Thr Ser  
 20 25 30  
 GGC TAT AGT TAT ATG CAC TGG AAC CAA CAG AAA CCA GGA CAG CCA CCC 144  
 Gly Tyr Ser Tyr Met His Trp Asn Gln Gln Lys Pro Gly Gln Pro Pro  
 35 40 45  
 AGA CTC CTC ATC TAT CTT GTA TCC AAC CTA GAA TCT GGG GTC CCT GCC 192  
 Arg Leu Leu Ile Tyr Leu Val Ser Asn Leu Glu Ser Gly Val Pro Ala  
 50 55 60  
 AGG TTC AGT GGC AGT GGG TCT GGG ACA GAC TTC ACC CTC AAC ATC CAT 240  
 Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Asn Ile His  
 65 70 75  
 CCT GTG GAG GAG GAG GAT GCT GCA ACC TAT TAC TGT CAG CAC ATT AGG 288  
 Pro Val Glu Glu Glu Asp Ala Ala Thr Tyr Tyr Cys Gln His Ile Arg  
 85 90 95  
 GGA GCT TAC ACG TTC GGA GGG GGG ACC AAG CTG GAA ATA AAA 330  
 Gly Ala Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys  
 100 105 110

(2) INFORMATION FOR SEQ ID NO: 37:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH:330 base pairs  
 (B) TYPE:nucleic acid  
 (C) STRANDEDNESS:double  
 (D) TOPOLOGY:linear  
 (ii) MOLECULE TYPE:mRNA  
 (iii) HYPOTHETICAL:no  
 (iv) ANTISENSE:no  
 (vi) ORIGINAL SOURCE:  
 (A) ORGANISM:mouse  
 (ix) FEATURE:  
 (A) NAME/KEY:Idio 20 L chain variable  
 (xi) SEQUENCE DESCRIPTION:SEQ ID NO: 37:

5 GAC ATT GTG CTG ACA CAG TCT CCT GCT TCC TTA GCT GTA TCT CTG GGG 48  
Asp Ile Val Leu Thr Gln Ser Pro Ala S r Leu Ala Val Ser Leu Gly 15

10 CAG AGG GCC ACC ATC TCA TAC AGG GCC AGC AAA AGT GTC AGT ACA TCT 96  
Gln Arg Ala Thr Il S r Tyr Arg Ala Ser Lys Ser Val Ser Thr Ser 20 25 30

15 GGC TAT AGT TAT ATG CAC TGG AAC CAA CAG AGA CCA GGA CAG CCA CCC 144  
Gly Tyr Ser Tyr Met His Trp Asn Gln Gln Arg Pro Gly Gln Pro Pro 35 40 45

20 AGA CTC CTC ATC TAT CTT GTA TCC AAC CTA GAC TCT GGG GTC CCT GCC 192  
Arg Leu Leu Ile Tyr Leu Val Ser Asn Leu Asp Ser Gly Val Pro Ala 50 55 60

25 AGG TTC AGT GGC AGT GGG TCT GGG ACA GAC TTC ACC CTC AAC ATC CAT 240  
Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Asn Ile His 65 70 75 80

30 CCT GTG GAG GAG GAG GAT GCT GCA ACC TAT TAC TGT CAG CAC ATT GAG 288  
Pro Val Glu Glu Glu Asp Ala Ala Thr Tyr Tyr Cys Gln His Ile Glu 85 90 95

35 GGA GCT TAC ACG TTC GGA GGG GGG ACC AAG CTG GAA ATA AAA 330  
Gly Ala Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys 100 105 110

## (2) INFORMATION FOR SEQ ID NO: 38:

30 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 321 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: double  
(D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: mRNA  
(iii) HYPOTHETICAL: no  
(iv) ANTISENSE: no  
(vi) ORIGINAL SOURCE:  
(A) ORGANISM: mouse

40 (ix) FEATURE:  
(A) NAME/KEY: Idio 27 L chain variable

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 38:

45 GAC ATT GTG ATG ACC CAG TCT CAC AAA TTC ATG TCC ACA TCA GTA GGA 48  
Asp Ile Val Met Thr Gln Ser His Lys Phe Met Ser Thr Ser Val Gly 5 10 15

50 GAC AGG GTC AGT ATC ACC TGC AAG GCC AGT CAG GAT GTG AAT ACT GCT 96  
Asp Arg Val Ser Ile Thr Cys Lys Ala Ser Gln Asp Val Asn Thr Ala 20 25 30

55 GTA GCC TGG TAT CAA CAG AAA CCA GGA CAA TCT CCT AAA CTA CTG CTT 144  
Val Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys Leu Leu Leu 35 40 45

TAC TCG GCA TCC TAC CGG TAC ACT GGA GTC CCT GAT CAC TTC ACT GGC 192  
Tyr Ser Ala Ser Tyr Arg Tyr Thr Gly Val Pro Asp His Phe Thr Gly  
50 55 60

AGT GGA TCT GGG ACG GAT TTC ACT TTC ACC ATC AGC GGT GTG CAG GCT 240  
Ser Gly Ser Gly Thr Asp Phe Thr Ph Thr Ile Ser Gly Val Gln Ala  
65 70 75 80

GAA GAC CTG GCA GTT TAT TAC TGT CAG CAA CAT TAT AGT CCT CCT CTC 288  
Glu Asp Leu Ala Val Tyr Tyr Cys Gln Gln His Tyr Ser Pro Pro Leu  
85 90 95

ACG TTC GGT GCT GGG ACC AAG CTG GAA CTG AAA 321  
Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys  
100 105

(2) INFORMATION FOR SEQ ID NO: 39:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 321 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: mRNA

(iii) HYPOTHETICAL: no

(iv) ANTISENSE: no

(vi) ORIGINAL SOURCE:

(A) ORGANISM: mouse

(ix) FEATURE:

(A) NAME/KEY: Idio 33 L chain variable

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 39:

GAC ATT GTG ATG ACA CAG TCT CAC AAA TTC ATG TCC ACA TCA GTT GGA 48  
Asp Ile Val Met Thr Gln Ser His Lys Phe Met Ser Thr Ser Val Gly  
5 10 15

GAC AGG GTC ACC ATC ACC TGC AAG GCC AGT CAG GAT GTG ACT ACT GAT 96  
Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Val Thr Thr Asp  
20 25 30

GTA GCC TGG TAT CAA CAG AAA CCA CGA CAA TCT CCT AAA CTA CTG ATT 144  
Val Ala Trp Tyr Gln Gln Lys Pro Arg Gln Ser Pro Lys Leu Leu Ile  
35 40 45

TAC TCG GCA TCC TAT CGG TAC ACT GGA GTC CCT GAT CGC TTC ACT GGC 192  
Tyr Ser Ala Ser Tyr Arg Tyr Thr Gly Val Pro Asp Arg Phe Thr Gly  
50 55 60

ACT GGA TCT GGG ACG GAT TTC ACT TTC ACC ATC AGC AGT GTG CAG GCT 240  
Ser Gly Ser Gly Thr Asp Phe Thr Phe Thr Ile Ser Ser Val Gln Ala  
65 70 75 80

GAA GAC CTG GCA GTT TAT TAC TGT CAG CAA CAT TAT AGT ACT GCG TGG 288  
Glu Asp Leu Ala Val Tyr Tyr Cys Gln Gln His Tyr Ser Thr Ala Trp  
85 90 95

ACG TTC GGT GGT GGC ACC AAG CTG GAA ATC AAA 321  
 Thr Phe Gly Gly Thr Lys Leu Glu Ile Lys  
 100 105

(2) INFORMATION FOR SEQ ID NO: 40:

(1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH:399 base pairs  
 (B) TYPE:nucleic acid  
 (C) STRANDEDNESS:double  
 (D) TOPOLOGY:linear

(ii) MOLECULE TYPE:mRNA

(iii) HYPOTHETICAL:no

(iv) ANTISENSE:no

(vi) ORIGINAL SOURCE:

(A) ORGANISM:mouse

(ix) FEATURE:

(A) NAME/KEY:Clone 3GB1

(xi) SEQUENCE DESCRIPTION:SEQ ID NO:40:

CTG TCG GTA ACT TCA GGG GTC TAC TCA GAG GTT CAG CTC GAG CAG TCT 48  
 Leu Ser Val Thr Ser Gly Val Tyr Ser Glu Val Gln Leu Gln Gln Ser  
 -5 1 5

GGG ACT GTG CTG GCA AGG CCT GGG GCT TCA GTG AAG ATG TCC TGC AAG 96  
 Gly Thr Val Leu Ala Arg Pro Gly Ala Ser Val Lys Met Ser Cys Lys  
 10 15 20

GCT TCG GGC TAC ACC TTT AAC AGC TAC TGG ATG CAC TGG GTA AAA CAG 144  
 Ala Ser Gly Tyr Thr Phe Asn Ser Tyr Trp Met His Trp Val Lys Gln  
 25 30 35

AGG CCT GGA CAG GGT CTG GAA TGG ATT GGC GCG ATT TAT CCT GGA AAT 192  
 Arg Pro Gly Gln Gly Leu Glu Trp Ile Gly Ala Ile Tyr Pro Gly Asn  
 40 45 50 55

AGT GAT ATT AGC TAC AGC CAG AAC TTT AAG GAC AGG GCC AAA CTG ACT 240  
 Ser Asp Ile Ser Tyr Ser Gln Asn Phe Lys Asp Arg Ala Lys Leu Thr  
 60 65 70

GCC GTC ACA TCC ACC AGC ACT GCC TAC ATG GAA CTC AGA AGC CTG ACA 288  
 Ala Val Thr Ser Thr Ser Thr Ala Tyr Met Glu Leu Arg Ser Leu Thr  
 75 80 85

AAT GAG GAC TCT GCG GTC TAT TTC TGT ACA AAA GAG GAA TAT GAT TAC 336  
 Asn Glu Asp Ser Ala Val Tyr Phe Cys Thr Lys Glu Glu Tyr Asp Tyr  
 90 95 100

GAC ACC CTG GAC TAC TGG GGT CAA GGA ACC TCA GTC ACC GTC TCC TCA 384  
 Asp Thr Leu Asp Tyr Trp Gly Gln Gly Thr Ser Val Thr Val Ser Ser  
 105 110 115

GCC AAA ACG ACA CCC 399  
 Ala Lys Thr Thr Pro  
 120

## (2) INFORMATION FOR SEQ ID NO: 41:

## (1) SEQUENCE CHARACTERISTICS:

- (A) LENGTH:402 base pairs  
 (B) TYPE:nucleic acid  
 (C) STRANDEDNESS:doubl  
 (D) TOPOLOGY:linear

(ii) MOLECULE TYPE:mRNA

(iii) HYPOTHETICAL:no

(iv) ANTISENSE:no

(vi) ORIGINAL SOURCE:

(A) ORGANISM:mouse

(ix) FEATURE:

(A) NAME/KEY:Clone 17GB7

(xi) SEQUENCE DESCRIPTION:SEQ ID NO: 41:

```

ATT GTG TCG GTA ACT TCA GGG GTC TAC TCA GAG GTT CAG CTC GAG CAG 48
Ile Leu Ser Val Thr Ser Gly Val Tyr Ser Glu Val Gln Leu Gln Gln
-10 -5 1 5

TCT GGG ACT GTG CTG GCA AGG CCT GGG GCT TCA GTG AAG ATG TCC TGC 96
Ser Gly Thr Val Leu Ala Arg Pro Gly Ala Ser Val Lys Met Ser Cys
10 15 20

AAG GCT TCG GGC TAC ACC TTT AAC AGC TAC TGG ATG CAC TGG GTA AAA 144
Lys Ala Ser Gly Tyr Thr Phe Asn Ser Tyr Trp Met His Trp Val Lys
25 30 35

CAG AGG CCT GGA CAG GGT CTG GAA TGG ATT GGC GCG ATT TAT CCT GGA 192
Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile Gly Ala Ile Tyr Pro Gly
40 45 50

AAT AGT GAT ATT AGC TAC AGC CAG AAC TTT AAG GAC AGG GCC AAA CTG 240
Asn Ser Asp Ile Ser Tyr Ser Gln Asn Phe Lys Asp Arg Ala Lys Leu
55 60 65 70

ACT GCC GTC ACA TCC ACC AGC ACT GAC TAC ATG GAA CTC AGA AGC CAG 288
Thr Ala Val Thr Ser Thr Ser Thr Ala Tyr Met Glu Leu Arg Ser Leu
75 80 85

ACA AAT GAG GAC TCT GCG GTC TAT TTC TGT ACA AAA GAG GAA TAT GAT 336
Thr Asn Glu Asp Ser Ala Val Tyr Phe Cys Thr Lys Glu Glu Tyr Asp
90 95 100

TAC GAC ACC CTG GAC TAC TGG GGT CAA GGA ACC TCA GTC ACC GTC TCC 384
Tyr Asp Thr Leu Asp Tyr Trp Gly Gln Gly Thr Ser Val Thr Val Ser
105 110 115

TCA GCC AAA ACG ACA CCC 402
Ser Ala Lys Thr Thr Pro
120

```

## (2) INFORMATION FOR SEQ ID NO: 42:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH:438 base pairs  
 (B) TYPE:nucleic acid

(C) STRANDEDNESS:double  
(D) TOPOLOGY:linear

5 (ii) MOLECULE TYPE:mRNA  
(iii) HYPOTHETICAL:no  
(iv) ANTISENSE:no  
(vi) ORIGINAL SOURCE:  
(A) ORGANISM:mouse  
(ix) FEATURE:  
10 (A) NAME/KEY:Clone 20GA2  
(xi) SEQUENCE DESCRIPTION:SEQ ID NO:42:

ATG GAG TTC GGG CTA AAC TGG GTT TTC CTT GTA ACA CTT TTA AAT GGT 48  
Met Glu Phe Gly Leu Asn Trp Val Phe Leu Val Thr Leu Leu Asn Gly  
-15 -10 -5

15 ATC CAG TGT GAG GTG AAG CTG GTG GAG TCT GGA GGA GGC TTG GTA CAG 96  
Ile Gln Cys Glu Val Lys Leu Val Glu Ser Gly Gly Gly Leu Val Gln  
1 5 10

20 CCT GGG GGT TCT CTC AGA CTC TCC TGT GCA ACT TCT GGG TTA ACC TTC 144  
Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Thr Ser Gly Leu Thr Phe  
15 20 25

25 ACT GAT TAC TAC ATG AAC TGG GTC CGC CAG CCT CCA GGA AAG GAA CTT 192  
Thr Asp Tyr Tyr Met Asn Trp Val Arg Gln Pro Pro Gly Lys Glu Leu  
30 35 40 45

GAA TGG TTG GGT TTT ATT AGA AAC AAA GCT AAT CTT TAC ACA ACA GAC 240  
Glu Trp Leu Gly Phe Ile Arg Asn Lys Ala Asn Leu Tyr Thr Thr Asp  
50 55 60

30 TAC AGT GCA TCT GTG AAG GGT CGG TTC ACC ATC TCC AGA CAT AAT CCC 288  
Tyr Ser Ala Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Pro  
65 70 75

35 CAA AGC ATC CTC TAT CTT CAA ATG AAC ACC CTG ACA ACT GAG GAC AGT 336  
Gln Ser Ile Leu Tyr Leu Gln Met Asn Thr Leu Thr Thr Glu Asp Ser  
80 85 90

GCC ACT TAT TAC TGT GCA AGA GAT AGG GGG GGG AGG GAC TGG TAC TTC 384  
Ala Thr Tyr Tyr Cys Ala Arg Asp Arg Gly Gly Arg Asp Trp Tyr Phe  
95 100 105

40 GAT GTC TGG GGC GCA GGG ACC ACG GTC ACC GTC TCC TCA GCC AAA ACG 432  
Asp Val Trp Gly Ala Gly Thr Thr Val Thr Val Ser Ser Ala Lys Thr  
110 115 120 125

45 ACA CCC 438  
Thr Pro

## (2) INFORMATION FOR SEQ ID NO: 43:

50 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH:411 base pairs  
(B) TYPE:nucleic acid  
(C) STRANDEDNESS:double  
(D) TOPOLOGY:linear

55

(ii) MOLECULE TYPE:mRNA  
 (iii) HYPOTHETICAL:no  
 (iv) ANTISENSE:no  
 (vi) ORIGINAL SOURCE:  
 (A) ORGANISM:mous  
 (ix) FEATURE:  
 (A) NAME/KEY:Clone 27GA5  
 (xi) SEQUENCE DESCRIPTION:SEQ ID NO: 43:

CTT GTA ACA CGT TTA AAT GGT ATC CAG TGT GAG GTG AAG CTG GTG GAG 48  
 Leu Val Thr Arg Leu Asn Gly Ile Gln Cys Glu Val Lys Leu Val Glu  
 -10 -5 1 5

TCT GGA GGA GGC TTG GTA CAG CCT GGG GGT TCT CTG AGA CTC TCC TGT 96  
 Ser Gly Gly Gly Leu Val Gln Pro Gly Gly Ser Leu Arg Leu Ser Cys  
 10 15 20

GCA ACT TCT GGG TTC ACC TTC ACT GAT TAC TAC ATG AAC TGG GTC CGC 144  
 Ala Thr Ser Gly Phe Thr Phe Thr Asp Tyr Tyr Met Asn Trp Val Arg  
 25 30 35

CAG CCT CCA GGA AAG GCA CTT GAG TGG TTG GGT TTT ATT AGA AAC AAA 192  
 Gln Pro Pro Gly Lys Ala Leu Glu Trp Leu Gly Phe Ile Arg Asn Lys  
 40 45 50

GCT AAT TAT TAC ACA ACA GAG TAC AGT GCA TCT GTG AAG GGT CGG TTC 240  
 Ala Asn Tyr Tyr Thr Thr Glu Tyr Ser Ala Ser Val Lys Gly Arg Phe  
 55 60 65 70

ACC ATC TCC AGA GAT AAT TCC CAA ACC ATC CTC TAT CTT CAA ATG AAC 288  
 Thr Ile Ser Arg Asp Asn Ser Gln Ser Ile Leu Gln Met Asn Thr Leu  
 75 80 85

ACC CTG AGA GCT GAG GAC AGT GCC ACT TAT TAC TGT GCA AGA GAT GGG 336  
 Thr Leu Arg Ala Glu Asp Ser Ala Thr Tyr Tyr Cys Ala Arg Asp Gly  
 90 95 100

TTC CTA CGG GAC TGG TAC TTC GAT GTC TGG GGC GCA GGG ACC ACG GTC 384  
 Phe Leu Arg Asp Trp Tyr Phe Asp Val Trp Gly Ala Gly Thr Thr Val  
 105 110 115

ACC GTC TCC TCA GCC AAA ACG ACA CCC GAT GAG GAG GAG GAG GAG 411  
 Thr Val Ser Ser Ala Lys Thr Thr Pro Gln Gln Gln Gln Gln Gln  
 120 125

(2) INFORMATION FOR SEQ ID NO: 44:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH:354 base pairs  
 (B) TYPE:nucleic acid  
 (C) STRANDEDNESS:double  
 (D) TOPOLOGY:linear

(ii) MOLECULE TYPE:mRNA  
 (iii) HYPOTHETICAL:no  
 (iv) ANTISENSE:no  
 (vi) ORIGINAL SOURCE:  
 (A) ORGANISM:mouse



## (ix) FEATURE:

(A) NAME/KEY:Clone 3KB11

## (xi) SEQUENCE DESCRIPTION:SEQ ID NO:44:

GAC ATT GTG CTG ACA CAG TCT CCT GCT TCC TTA GCT GTA TCT CCT CTG 48  
 Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Pro Leu  
 5 10 15

GGG CAG AGG GCC ACC ATC TCA TAC AGG GCC AGC AAA AGT GTG CAG TTA 96  
 Gly Gln Arg Ala Thr Ile Ser Tyr Arg Ala Ser Lys Ser Val Gln Leu  
 20 25 30

CAT CTG GCT ATA GTT TAT ATG CAC TGG AAC CAA CAG AAA CCA GGA CAG 144  
 His Leu Ala Ile Val Tyr Met His Trp Asn Gln Gln Lys Pro Gly Gln  
 35 40 45

CCA CCC AGA CTC CTC ATC TAT CTT GTA TCC AAC CTA GAA TCT GGG GTC 192  
 Pro Pro Arg Leu Leu Ile Tyr Leu Val Ser Asn Leu Glu Ser Gly Val  
 50 55 60

CCT GCC AGG TTC AGT GGC AGT GGG TCT GGG ACA GAC TTC ACC CTC AAC 240  
 Pro Ala Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Asn  
 65 70 75 80

ATC CAT CCT GTG GAG GAG GAG GAT GCT GCA ACC TAT TAC TGT CAG CAC 288  
 Ile His Pro Val Glu Glu Glu Asp Ala Ala Thr Tyr Tyr Cys Gln His  
 85 90 95

ATT AGG GTA GCT TAC ACG TTC GGA GGG GGG ACC AAG CTG GAA ATA AAA 336  
 Ile Arg Val Ala Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys  
 100 105 110

CGG GCT GAT GCT GCA CCA GAG TAT TCA TCA TCA TCA TCA TCA TCA TCA 354  
 Arg Ala Asp Ala Ala Pro Tyr Tyr Tyr Tyr Tyr Tyr Tyr Tyr Tyr  
 115 120 125 130 135 140 145 150

## (2) INFORMATION FOR SEQ ID NO: 45:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH:438 base pairs

(B) TYPE:nucleic acid

(C) STANDEDNESS:double

(D) TOPOLOGY:linear

(ii) MOLECULE TYPE:mRNA

(iii) HYPOTHETICAL:no

(iv) ANTISENSE:no

(vi) ORIGINAL SOURCE:

(A) ORGANISM:mouse

## (ix) FEATURE:

(A) NAME/KEY:Clone 17KB1

## (xi) SEQUENCE DESCRIPTION:SEQ ID NO: 45:

CTA TGG GTA CTG CTG CTC TGG GTT CCA GGT TCC ACT GGT GAC ATT GTG 48  
 Leu Trp Val Leu Leu Leu Trp Val Pro Gly Ser Thr Gly Asp Ile Val  
 -10 -5

5 CTG ACA CAG TCT CCT GCT TCC TTA GCT GTA TCT CTG GGG CAG AGG GCC 96  
 Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Leu Gly Gln Arg Ala  
 5 10 15  
 TCC ATC TCA TAC AGG GCC AGC AAA AGT GTC AGT ACA TCT GGC TAT AGT 144  
 Ser Ile Ser Tyr Arg Ala Ser Lys Ser Val Ser Thr Ser Gly Tyr Ser  
 20 25 30 35  
 10 TAT ATG CAC TGG AAC CAA CAG AAA CCA GGA CAG CCA CCC AGA CTC CTC 192  
 Tyr Met His Trp Asn Gln Gln Lys Pro Gly Gln Pro Pro Arg Leu Leu  
 40 45 50  
 ATC TAT CTT GTA TCC AAC CTA GAA TCT GGG GTC CCT GCC AGG TTC AGT 240  
 Ile Tyr Leu Val Ser Asn Leu Glu Ser Gly Val Pro Ala Arg Phe Ser  
 15 55 60 65  
 GGC AGT GGG TCT GGG ACA GAC TTC ACC CTC AAC ATC CAT CCT GTG GAG 288  
 Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Asn Ile His Pro Val Glu  
 70 75 80  
 20 GAG GAG GAT GCT GCA ACC TAT TAC TGT CAG CAC ATT AGG GGA GCT TAC 336  
 Glu Glu Asp Ala Ala Thr Tyr Tyr Cys Gln His Ile Arg Gly Ala Tyr  
 85 90 95  
 ACG TTC GGA GGG GGG ACC AAG CTG GAA ATA AAA CGG GCT GAT GCT GCA 394  
 Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg Ala Asp Ala Ala  
 100 105 110 115  
 CCA ACT GTA TCC ATC TTC CCA CCA TCC AGT AAG CTT GGG AAA CGG TTC 432  
 Pro Thr Val Ser Ile Phe Pro Pro Ser Ser Lys Leu Gly Lys Arg Phe  
 120 125 130  
 30 GCA CCG 438  
 Ala Pro

## (2) INFORMATION FOR SEQ ID NO: 46:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 417 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: mRNA

(iii) HYPOTHETICAL: no

(iv) ANTISENSE: no

(vi) ORIGINAL SOURCE:

(A) ORGANISM: mouse

(ix) FEATURE:

(A) NAME/KEY: Clone 20KB1

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 46:

GGCCGCG GTGAGAACCG TTGGGAATTC ATG GAG ACA GAC ACA CTC CTC 48  
 Met Glu Thr Asp Thr Leu Leu  
 -20 -15

5 CTA TGG GTA CTG CTG CTC TGG GTT CCA GGT TCC ACT GGT GAC ATT GTG 96  
 Leu Trp Val Leu Leu Trp Val Pro Gly Ser Thr Gly Asp Ile Val  
 -10 -5 1

CTG ACA CAG TCT CCT GCT TCC TTA GCT GTA TCT CTG GGG CAG AGG GCC 144  
 Leu Thr Gln Ser Pro Ala Ser Leu Ala Val S r Leu Gly Gln Arg Ala  
 5 10 15

10 ACC ATC TCA TAC AGG GCC AGC AAA AGT GTC AGT ACA TCT GGC TAT AGT 192  
 Thr Ile Ser Tyr Arg Ala Ser Lys Ser Val Ser Thr Ser Gly Tyr Ser  
 20 25 30

15 TAT ATG CAC TGG AAC CAA CAG AGA CCA GGA CAG CCA CCC AGA CTC CTC 240  
 Tyr Met His Trp Asn Gln Gln Arg Pro Gly Gln Pro Pro Arg Leu Leu  
 35 40 45

ATC TAT CTT GTA TCC AAC CTA GAC TCT GGG GTC CCT GCC AGG TTC AGT 288  
 Ile Tyr Leu Val Ser Asn Leu Asp Ser Gly Val Pro Ala Arg Phe Ser  
 50 55 60 65

20 GGC AGT GGG TCT GGG ACA GAC TTC ACC CTC AAC ATC CAT CCT GTG GAG 336  
 Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Asn Ile His Pro Val Glu  
 70 75 80

25 GAG GAG GAT GCT GCA ACC TAT TAC TGT CAG CAC ATT GAG GGA GCT TAC 384  
 Glu Glu Asp Ala Ala Thr Tyr Tyr Cys Gln His Ile Glu Gly Ala Tyr  
 85 90 95

ACG TTC GGA GGG GGG ACC AAG CTG GAA ATA AAA 417  
 Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys  
 100 105

## (2) INFORMATION FOR SEQ ID NO: 47:

35 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH:420 base pairs  
 (B) TYPE:nucleic acid  
 (C) STRANDEDNESS:single  
 (D) TOPOLOGY:linear

(ii) MOLECULE TYPE:mRNA

40 (iii) HYPOTHETICAL:no

(iv) ANTISENSE:no

(vi) ORIGINAL SOURCE:  
 (A) ORGANISM:mouse

(ix) FEATURE:  
 (A) NAME/KEY:Clone 27KA2

45 (xi) SEQUENCE DESCRIPTION:SEQ ID NO: 47:  
 GCGGCCGCGG TGAGAACCGT TTGGGAATTC ATC GAG ACA CAG TCC CAG 48  
 Met Glu Thr Gln Ser Gln  
 -20 -15

50 GTC TTT GTA TTC GTG TTT CTC TGG TTG TCT GGT GTT GAC GGA GAC ATT 96  
 Val Phe Val Phe Val Phe Leu Trp Leu Ser Gly Val Asp Gly Asp Ile  
 -10 -5 1

GTG ATG ACC CAG TCT CAC AAA TTC ATG TCC ACA TCA GTA GGA GAC AGG 144  
Val Met Thr Gln Ser His Lys Phe Met Ser Thr Ser Val Gly Asp Arg  
5 5 10 15

GTC AGT ATC ACC TGC AAG GCC AGT CAG GAT GTG AAT ACT GCT GTA GCC 192  
Val Ser Ile Thr Cys Lys Ala Ser Gln Asp Val Asn Thr Ala Val Ala  
20 25 30

TGG TAT CAA CAG AAA CCA GGA CAA TCT CCT AAA CTA CTG CTT TAC TCG 240  
Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys Leu Leu Leu Tyr Ser  
35 40 45 50

GCA TCC TAC CGG TAC ACT GGA GTC CCT GAT CAC TTC ACT GGC AGT GGA 288  
Ala Ser Tyr Arg Tyr Thr Gly Val Pro Asp His Phe Thr Gly Ser Gly  
55 60 65

TCT GGG ACG GAT TTC ACT TTC ACC ATC AGC GGT GTG CAG GCT GAA GAC 336  
Ser Gly Thr Asp Phe Thr Phe Thr Ile Ser Gly Val Gln Ala Glu Asp  
70 75 80

CTG GCA GTT TAT TAC TGT CAG CAA CAT TAT AGT CCT CCT CTC ACG TTC 384  
Leu Ala Val Tyr Tyr Cys Gln Gln His Tyr Ser Pro Pro Leu Thr Phe  
85 90 95

GGT GCT GGG ACC AAG CTG GAA CTG AAA CGG GCT GAT 420  
Gly Ala Gly Thr Lys Leu Glu Leu Lys Arg Ala Asp  
100 105 110

(2) INFORMATION FOR SEQ ID NO: 48:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 360 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: mRNA

(iii) HYPOTHETICAL: no

(iv) ANTISENSE: no

(vi) ORIGINAL SOURCE:

(A) ORGANISM: mouse

(ix) FEATURE:

(A) NAME/KEY: Clone 23KA26

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 48

GGT GTT GAC GGA GAC ATT GTG ATG ACA CAG TCT CAC AAA TTC ATG TCC 48  
Gly Val Asp Gly Asp Ile Val Met Thr Gln Ser His Lys Phe Met Ser  
1 5 10

ACA TCA GTT GGA GAC AGG GTC ACC ATC ACC TGC AAG GCC AGT CAG GAT 96  
Thr Ser Val Gly Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp  
15 20 25

GTG ACT ACT GAT GTA GCC TGG TAT CAA CAG AAA CCA CGA CAA TCT CCT 144  
Val Thr Thr Asp Val Ala Trp Tyr Gln Gln Lys Pro Arg Gln Ser Pro  
30 35 40

AAA CTA CTG ATT TAC TCG GCA TCC TAT CGG TAC ACT GGA GTC CCT GAT 192  
 Lys Leu Leu Ile Tyr Ser Ala Ser Tyr Arg Tyr Thr Gly Val Pro Asp 195  
 5 45 50 55  
 CGC TTC ACT GGC AGT GGA TCT GGG ACG GAT TTC ACT TTC ACC ATC AGC 240  
 Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Ph Thr Phe Thr Ile Ser 245  
 60 65 70 75  
 10 AGT GTG CAG GCT GAA GAC CTG GCA GTT TAT TAC TGT CAG CAA CAT TAT 286  
 Ser Val Gln Ala Glu Asp Leu Ala Val Tyr Tyr Cys Gln Gln His Tyr 290  
 80 85 90  
 15 AGT ACT GCG TGG ACG TTC GGT GGT GGC ACC AAG CTG GAA ATC AAA CCG 336  
 Ser Thr Ala Trp Thr Phe Gly Gly Thr Lys Leu Glu Ile Lys Arg 340  
 95 100 105  
 GCT GAT GCT GCA CGA ACT GTA TCC 360  
 Ala Asp Ala Ala Pro Thr Val Ser 365  
 110 115

20 TAT TAT TAT TAT TAT TAT TAT TAT TAT TAT TAT TAT TAT TAT TAT TAT  
 TAT TAT TAT TAT TAT TAT TAT TAT TAT TAT TAT TAT TAT TAT TAT TAT  
 TAT TAT TAT TAT TAT TAT TAT TAT TAT TAT TAT TAT TAT TAT TAT TAT

## 25 Claims

1. An immunoglobulin H chain variable region fragment which contains a hypervariable region CDR1 having an amino acid sequence selected from

30 (1) Ser Tyr Trp Met His;  
 Asp Tyr Tyr Met Asn; and  
 Asn Tyr Trp Met Gln

35 a hypervariable region CDR2 having an amino acid sequence selected from

40 (2) Ala Ile Tyr Pro Gly Asn Ser  
 Asp Ile Ser Tyr Ser Gln Asn  
 Phe Lys Asp;  
 Phe Ile Arg Asn Lys Ala  
 45 Asn Leu Tyr Thr Thr Asp  
 Tyr Ser Ala Ser Val Lys  
 Gly;  
 Phe Ile Arg Asn Lys Ala  
 50 Asn Tyr Tyr Thr Thr Glu  
 Tyr Ser Ala Ser Val Lys  
 Gly; and  
 Ala Ile Tyr Pro Gly Asp  
 Gly Asp Thr Arg Tyr Thr  
 Gln Lys Phe Lys Gly,

and a hypervariable region CDHs having an amino acid sequence selected from

(3) Glu Glu Tyr Asp Tyr Asp  
 Thr Leu Asp Tyr;  
 Asp Arg Gly Gly Arg Asp  
 Trp Tyr Phe Asp Val;  
 Asp Gly Phe Leu Arg Asp  
 Trp Tyr Phe Asp Val; and  
 Ser Gly Tyr Tyr Gly Ser  
 Phe Val Gly Phe Ala Tyr .

2. An immunoglobulin H chain variable region fragment having the following amino acid sequence

Glu Val Gln Leu Gln Gln Ser Gly Thr Val  
 Leu Ala Arg Pro Gly Ala Ser Val Lys Met  
 Ser Cys Lys Ala Ser Gly Tyr Thr Phe Asn  
 Ser Tyr Trp Met His Trp Val Lys Gln Arg  
 Pro Gly Gln Gly Leu Glu Trp Ile Gly Ala  
 Ile Tyr Pro Gly Asn Ser Asp Ile Ser Tyr  
 Ser Gln Asn Phe Lys Asp Arg Ala Lys Leu  
 Thr Ala Val Thr Ser Thr Ser Thr Ala Tyr  
 Met Glu Leu Arg Ser Leu Thr Asn Glu Asp  
 Ser Ala Val Tyr Phe Cys Thr Lys Glu Glu  
 Tyr Asp Tyr Asp Thr Leu Asp Tyr Trp Gly  
 Gln Gly Thr Ser Val Thr Val Ser Ser

## 3. An immunoglobulin H chain variable region fragment having the following amino acid sequence

5      Glu Val Lys Leu Val Glu Ser Gly Gly Gly  
       Leu Val Gln Pro Gly Gly Ser Leu Arg Leu  
       Ser Cys Ala Thr Ser Gly Phe Thr Phe Thr  
       Asp Tyr Tyr Met Asn Trp Val Arg Gln Pro  
 10     Pro Gly Lys Ala Leu Glu Trp Leu Gly Phe  
       Ile Arg Asn Lys Ala Asn Tyr Tyr Thr Thr  
       Glu Tyr Ser Ala Ser Val Lys Gly Arg Phe  
       Thr Ile Ser Arg Asp Asn Ser Gln Ser Ile  
 15     Leu Tyr Leu Gln Met Asn Thr Leu Arg Ala  
       Glu Asp Ser Ala Thr Tyr Tyr Cys Ala Arg  
       Asp Gly Phe Leu Arg Asp Trp Tyr Phe Asp  
       Val Trp Gly Ala Gly Thr Thr Val Thr Val  
 20     Ser Ser.

## 25 4. An immunoglobulin H chain variable region fragment having the following amino acid sequence

30     Glu Val Lys Leu Val Glu Ser Gly Gly Gly  
       Leu Val Gln Pro Gly Gly Ser Leu Arg Leu  
       Ser Cys Ala Thr Ser Gly Leu Thr Phe Thr  
       Asp Tyr Tyr Met Asn Trp Val Arg Gln Pro  
       Pro Gly Lys Glu Leu Glu Trp Leu Gly Phe  
 35     Ile Arg Asn Lys Ala Asn Leu Tyr Thr Thr  
       Asp Tyr Ser Ala Ser Val Lys Gly Arg Phe  
       Thr Ile Ser Arg Asp Asn Pro Gln Ser Ile  
       Leu Tyr Leu Gln Met Asn Thr Leu Thr Thr  
 40     Glu Asp Ser Ala Thr Tyr Tyr Cys Ala Arg  
       Asp Arg Gly Gly Arg Asp Trp Tyr Phe Asp  
       Val Trp Gly Ala Gly Thr Thr Val Thr Val  
 45     Ser Ser.

5. An immunoglobulin H chain variable region fragment having the following amino acid sequence:

Glu Val Gln Leu Gln Gln Ser Gly Ala Glu  
 Leu Ala Arg Pro Gly Ala Ser Val Asn Leu  
 Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr  
 Asn Tyr Trp Met Gln Trp Val Lys Gln Arg  
 Pro Gly Gln Gly Leu Glu Trp Ile Gly Ala  
 Ile Tyr Pro Gly Asp Gly Asp Thr Arg Tyr  
 Thr Gln Lys Phe Lys Gly Lys Ala Thr Leu  
 Thr Ala Ala Lys Ser Ser Ser Thr Ala Tyr  
 Met Gln Leu Ser Ser Leu Ala Ser Glu Asp  
 Ser Ala Val Tyr Tyr Cys Ala Arg Ser Gly  
 Tyr Tyr Gly Ser Phe Val Gly Phe Ala Tyr  
 Trp Gly Gln Gly Thr Leu Val Thr Val Ser  
 Ala .

6. DNA and RNA fragments each encoding an immunoglobulin H chain variable region fragment which contains a base sequence encoding a hypervariable region CDR1 having an amino acid sequence selected from

(1) Ser Tyr Trp Met His;

Asp Tyr Tyr Met Asn; and

Asn Tyr Trp Met Gln;

a base sequence encoding a hypervariable region CDR2 having an amino acid sequence selected from

(2) Ala Ile Tyr Pro Gly Asn Ser

Asp Ile Ser Tyr Ser Gln Asn

Phe Lys Asp;

Phe Ile Arg Asn Lys Ala

Asn Leu Tyr Thr Thr Asp

Tyr Ser Ala Ser Val Lys

Gly;

Phe Ile Arg Asn Lys Ala

Asn Tyr Tyr Thr Thr Glu

Tyr Ser Ala Ser Val Lys

Gly; and

Ala Ile Tyr Pro Gly Asp

Gly Asp Thr Arg Tyr Thr

Glu Lys Phe Lys Gly .



a base sequence encoding a hypervariable region CDR3 having an amino acid sequence selected from

(3) Glu Glu Tyr Asp Tyr Asp  
 Thr Leu Asp Tyr;  
 Asp Arg Gly Gly Arg Asp  
 Trp Tyr Phe Asp Val;  
 Asp Gly Phe Leu Arg Asp  
 Trp Tyr Phe Asp Val; and  
 Ser Gly Tyr Tyr Gly Ser  
 Phe Val Gly Phe Ala Tyr.

7. An immunoglobulin H chain variable region fragment having following base sequence:

GAG GTT CAG CTC CAG CAG TCT GGG ACT GTG  
 CTG GCA AGG CCT GGG GCT TCA GTG AAG ATG  
 TCC TGC AAG GCT TCG GGC TAC ACC TTT AAC  
 AGC TAC TGG ATG CAC TGG GTA AAA CAG AGG  
 CCT GGA CAG GGT CTG GAA TGG ATT GGC GCG  
 ATT TAT CCT GGA AAT AGT GAT ATT AGC TAC  
 AGC CAG AAC TTT AAG GAC AGG GCC AAA CTG  
 ACT GCC GTC ACA TCC ACC AGC ACT GCC TAC  
 ATG GAA CTC AGA AGC CTG ACA AAT GAG GAC  
 TCT GCG GTC TAT TTC TGT ACA AAA GAG GAA  
 TAT GAT TAC GAC ACC CTG GAC TAC TGG GGT  
 CAA GGA ACC TCA GTC ACC GTC TCC TCA.

8. An immunoglobulin H chain variable region fragment having the following base sequence.

5 GAG GTG AAG CTG GTG GAG TCT GGA GGA GGC  
 TTG GTA CAG CCT GGG GGT TCT CTC AGA CTC  
 TCC TGT GCA ACT TCT GGG TTA ACC TTC ACT  
 GAT TAC TAC ATG AAC TGG GTC CGC CAG CCT  
 10 CCA GGA AAG GAA CTT GAA TGG TTG GGT TTT  
 ATT AGA AAC AAA GCT AAT CTT TAC ACA ACA  
 GAC TAC AGT GCA TCT GTG AAG GGT CGG TTC  
 ACC ATC TCC AGA GAT AAT CCC CAA AGC ATC  
 15 CTC TAT CTT CAA ATG AAC ACC CTG ACA ACT  
 GAG GAC AGT GCC ACT TAT TAC TGT GCA AGA  
 GAT AGG GGG GGG AGG GAC TGG TAC TTC GAT  
 20 GTC TGG GGC GCA GGG ACC ACG GTC ACC GTC  
 TCC TCA .

- 25 9. An immunoglobulin H chain variable region fragment having the following base sequence

30 GAG GTG AAG CTG GTG GAG TCT GGA GGA GGC  
 TTG GTA CAG CCT GGG GGT TCT CTG AGA CTC  
 TCC TGT GCA ACT TCT GGG TTC ACC TTC ACT  
 GAT TAC TAC ATG AAC TGG GTC CGC CAG CCT  
 CCA GGA AAG GCA CTT GAG TGG TTG GGT TTT  
 35 ATT AGA AAC AAA GCT AAT TAT TAC ACA ACA  
 GAG TAC AGT GCA TCT GTG AAG GGT CGG TTC  
 ACC ATC TCC AGA GAT AAT TCC CAA AGC ATC  
 CTC TAT CTT CAA ATG AAC ACC CTG AGA GCT  
 40 GAG GAC AGT GCC ACT TAT TAC TGT GCA AGA  
 GAT GGG TTC CTA GGG GAC TGG TAC TTC GAT  
 GTC TGG GGC GCA GGG ACC ACG GTC ACC GTC  
 45 TCC TCA .

## 10. An immunoglobulin H chain variable region fragment having the following base sequence

5 GAG GTT CAG CTC CAG CAG TCT GGG GCT GAA  
 CTG GCA AGA CCT GGG GCT TCA GTG AAC TTG  
 TCC TGC AAG GCT TCT GGC TAC ACC TTT ACT  
 AAC TAC TGG ATG CAG TGG GTA AAA CAG AGG  
 10 CCT GGA CAG GGT CTG GAA TGG ATT GGG GCT  
 ATT TAT CCT GGA GAT GGT GAT ACT AGG TAC  
 ACT CAG AAG TTC AAG GGC AAG GCC ACA TTG  
 ACT GCA GCT AAA TCC TCC AGC ACA GCC TAC  
 15 ATG CAA CTC AGC AGC TTG GCA TCT GAG GAC  
 TCT GCG GTC TAT TAC TGT GCA AGA TCG GGC  
 TAC TAT GGT AGC TTC GTT GGG TTT GCT TAC  
 TGG GGC CAA GGG ACT CTG GTC ACT GTC TCT  
 20 GCA .

## 25 11. An immunoglobulin L chain variable region fragment which contains a hypervariable region CDR1 having an amino acid sequence selected from

(1) Tyr Arg Ala Ser Lys Ser Val  
 30 Gln Leu His Leu Ala Ile Val  
 Tyr Met His  
 Tyr Arg Ala Ser Lys Ser Val  
 Ser Thr Ser Gly Tyr Ser Tyr  
 35 Met His  
 Lys Ala Ser Gln Asp Val Asn  
 Thr Ala Val Ala; and  
 40 Lys Ala Ser Gln Asp Val Thr  
 Thr Asp Val Ala

a hypervariable region CDR2 having an amino acid sequence selected from

(2) Leu Val Ser Asn Leu Glu Ser;  
 Leu Val Ser Asn Leu Asp Ser; and  
 50 Ser Ala Ser Tyr Arg Tyr Thr,

and a hypervariable region CDR3 having an amino acid sequence selected from:

(3) Gln His Ile Arg Val Ala Tyr  
 Thr;  
 Gln His Ile Arg Gly Ala Tyr  
 Thr;  
 Gln His Ile Glu Gly Ala Tyr  
 Thr;  
 Gln Gln His Tyr Ser Pro Pro  
 Leu Thr; and  
 Gln Gln His Tyr Ser Thr Ala  
 Trp Thr.

12. An immunoglobulin L chain variable region fragment having the following amino acid sequence

Asp Ile Val Leu Thr Gln Ser Pro Ala Ser  
 Leu Ala Val Ser Pro Leu Gly Gln Arg Ala  
 Thr Ile Ser Tyr Arg Ala Ser Lys Ser Val  
 Gln Leu His Leu Ala Ile Val Tyr Met His  
 Trp Asn Gln Gln Lys Pro Gly Gln Pro Pro  
 Arg Leu Leu Ile Tyr Leu Val Ser Asn Leu  
 Glu Ser Gly Val Pro Ala Arg Phe Ser Gly  
 Ser Gly Ser Gly Thr Asp Phe Thr Leu Asn  
 Ile His Pro Val Glu Glu Glu Asp Ala Ala  
 Thr Tyr Tyr Cys Gln His Ile Arg Val Ala  
 Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu  
 Ile Lys.

## 13. An immunoglobulin L chain variable region fragment having the following amino acid sequence

5      Asp Ile Val Leu Thr Gln Ser Pro Ala Ser  
       Leu Ala Val Ser Leu Gly Gln Arg Ala Ser  
       Ile Ser Tyr Arg Ala Ser Lys Ser Val Ser  
       Thr Ser Gly Tyr Ser Tyr Met His Trp Asn  
 10     Gln Gln Lys Pro Gly Gln Pro Pro Arg Leu  
       Leu Ile Tyr Leu Val Ser Asn Leu Glu Ser  
       Gly Val Pro Ala Arg Phe Ser Gly Ser Gly  
       Ser Gly Thr Asp Phe Thr Leu Asn Ile His  
 15     Pro Val Glu Glu Glu Asp Ala Ala Thr Tyr  
       Tyr Cys Gln His Ile Arg Gly Ala Tyr Thr  
       Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys.

20

## 14. An immunoglobulin L chain variable region fragment having the following amino acid sequence

25     Asp Ile Val Leu Thr Gln Ser Pro Ala Ser  
       Leu Ala Val Ser Leu Gly Gln Arg Ala Thr  
       Ile Ser Tyr Arg Ala Ser Lys Ser Val Ser  
 30     Thr Ser Gly Tyr Ser Tyr Met His Trp Asn  
       Gln Gln Arg Pro Gly Gln Pro Pro Arg Leu  
       Leu Ile Tyr Leu Val Ser Asn Leu Asp Ser  
       Gly Val Pro Ala Arg Phe Ser Gly Ser Gly  
 35     Ser Gly Thr Asp Phe Thr Leu Asn Ile His  
       Pro Val Glu Glu Glu Asp Ala Ala Thr Tyr  
       Tyr Cys Gln His Ile Glu Gly Ala Tyr Thr  
 40     Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys.

45

50

55

## 15. An immunoglobulin L chain variable region fragment having the following amino acid sequence

Asp Ile Val Met Thr Gln Ser His Lys Phe  
 Met Ser Thr Ser Val Gly Asp Arg Val Ser  
 Ile Thr Cys Lys Ala Ser Gln Asp Val Asn  
 Thr Ala Val Ala Trp Tyr Gln Gln Lys Pro  
 Gly Gln Ser Pro Lys Leu Leu Leu Tyr Ser  
 Ala Ser Tyr Arg Tyr Thr Gly Val Pro Asp  
 His Phe Thr Gly Ser Gly Ser Gly Thr Asp  
 Phe Thr Phe Thr Ile Ser Gly Val Gln Ala  
 Glu Asp Leu Ala Val Tyr Tyr Cys Gln Gln  
 His Tyr Ser Pro Pro Leu Thr Phe Gly Ala  
 Gly Thr Lys Leu Glu Leu Lys .

## 16. An immunoglobulin L chain variable region fragment having the following amino acid sequence

Asp Ile Val Met Thr Gln Ser His Lys Phe  
 Met Ser Thr Ser Val Gly Asp Arg Val Thr  
 Ile Thr Cys Lys Ala Ser Gln Asp Val Thr  
 Thr Asp Val Ala Trp Tyr Gln Gln Lys Pro  
 Arg Gln Ser Pro Lys Leu Leu Ile Tyr Ser  
 Ala Ser Tyr Arg Tyr Thr Gly Val Pro Asp  
 Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp  
 Phe Thr Phe Thr Ile Ser Ser Val Gln Ala  
 Glu Asp Leu Ala Val Tyr Tyr Cys Gln Gln  
 His Tyr Ser Thr Ala Trp Thr Phe Gly Gly  
 Gly Thr Lys Leu Glu Ile Lys .

17. DNA and RNA fragments each encoding an immunoglobulin L chain variable region fragment which contains a base sequence encoding a hypervariable region CDR1 having an amino acid sequence selected from

(1) Tyr Arg Ala Ser Lys Ser Val  
Gln Leu His Leu Ala Ile Val  
Tyr Met His;  
Tyr Arg Ala Ser Lys Ser Val  
Ser Thr Ser Gly Tyr Ser Tyr  
Met His;  
Lys Ala Ser Gln Asp Val Asn  
Thr Ala Val Ala; and  
Lys Ala Ser Gln Asp Val Thr  
Thr Asp Val Ala.

a base sequence encoding a hypervariable region CDR2 having an amino acid sequence selected from

(2) Leu Val Ser Asn Leu Glu Ser;  
Leu Val Ser Asn Leu Asp Ser; and  
Ser Ala Ser Tyr Arg Tyr Thr

and a base sequence encoding a hypervariable region CDR3 having an amino acid sequence selected from

(3) Gln His Ile Arg Val Ala Tyr  
Thr;  
Gln His Ile Arg Gly Ala Tyr  
Thr;  
Gln His Ile Glu Gly Ala Tyr  
Thr;  
Gln Gln His Tyr Ser Pro Pro  
Leu Thr; and  
Gln Gln His Tyr Ser Thr Ala  
Trp Thr .

## 18. An immunoglobulin L chain variable region fragment having the following base sequence

5 GAC ATT GTG CTG ACA CAG TCT CCT GCT TCC  
 TTA GCT GTA TCT CCT CTG GGG CAG AGG GCC  
 ACC ATC TCA TAC AGG GCC AGC AAA AGT GTG  
 CAG TTA CAT CTG GCT ATA GTT TAT ATG CAC  
 10 TGG AAC CAA CAG AAA CCA GGA CAG CCA CCC  
 AGA CTC CTC ATC TAT CTT GTA TCC AAC CTA  
 GAA TCT GGG GTC CCT GCC AGG TTC AGT GGC  
 AGT GGG TCT GGG ACA GAC TTC ACC CTC AAC  
 15 ATC CAT CCT GTG GAG GAG GAG GAT GCT GCA  
 ACC TAT TAC TGT CAG CAC ATT AGG GTA GCT  
 TAC ACG TTC GGA GGG GGG ACC AAG CTG GAA  
 20 ATA AAA .

## 19. An immunoglobulin L chain variable region fragment having the following base sequence

25 GAC ATT GTG CTG ACA CAG TCT CCT GCT TCC  
 TTA GCT GTA TCT CTG GGG CAG AGG GCC TCC  
 30 ATC TCA TAC AGG GCC AGC AAA AGT GTC AGT  
 ACA TCT GGC TAT AGT TAT ATG CAC TGG AAC  
 CAA CAG AAA CCA GGA CAG CCA CCC AGA CTC  
 CTC ATC TAT CTT GTA TCC AAC CTA GAA TCT  
 35 GGG GTC CCT GCC AGG TTC AGT GGC AGT GGG  
 TCT GGG ACA GAC TTC ACC CTC AAC ATC CAT  
 CCT GTG GAG GAG GAG GAT GCT GCA ACC TAT  
 40 TAC TGT CAG CAC ATT AGG GGA GCT TAC ACG  
 TTC GGA GGG GGG ACC AAG CTG GAA ATA AAA .



20. An immunoglobulin L chain variable region fragment having the following base sequence

5  
GAC ATT GTG CTG ACA CAG TCT CCT GCT TCC  
TTA GCT GTA TCT CTG GGG CAG AGG GCC ACC  
ATC TCA TAC AGG GCC AGC AAA AGT GTC AGT  
ACA TCT GGC TAT AGT TAT ATG CAC TGG AAC  
CAA CAG AGA CCA GGA CAG CCA CCC AGA CTC  
10 CTC ATC TAT CTT GTA TCC AAC CTA GAC TCT  
GGG GTC CCT GCC AGG TTC AGT GGC AGT GGG  
TGT GGG ACA GAC TTC ACC CTC AAC ATC CAT  
CCT GTG GAG GAG GAG GAT GCT GCA ACC TAT  
15 TAC TGT CAG CAC ATT GAG GGA GCT TAC ACG  
TTC GGA GGG GGG ACC AAG CTG GAA ATA AAA

20

21. An immunoglobulin L chain variable region fragment having the following base sequence

25 GAC ATT GTG ATG ACC CAG TCT CAC AAA TTC  
ATG TCC ACA TCA GTA GGA GAC AGG GTC AGT  
ATC ACC TGC AAG GCC AGT CAG GAT GTG AAT  
ACT GCT GTA GCC TGG TAT CAA CAG AAA CCA  
30 GGA CAA TCT CCT AAA CTA CTG CTT TAC TCG  
GCA TCC TAC CGG TAC ACT GGA GTC CCT GAT  
CAC TTC ACT GGC AGT GGA TCT GGG ACG GAT  
TTC ACT TTC ACC ATC AGC GGT GTG CAG GCT  
35 GAA GAC CTG GCA GTT TAT TAC TGT CAG CAA  
CAT TAT AGT CCT CCT CTC ACG TTC GGT GCT  
GGG ACC AAG CTG GAA CTG AAA

40

45

50

55

22. An immunoglobulin L chain variable region fragment having the following base sequence

5 GAC ATT GTG ATG ACA CAG TCT CAC AAA TTC  
 ATG TCC ACA TCA GTT GGA GAC AGG GTC ACC  
 ATC ACC TGC AAG GCC AGT CAG GAT GTG ACT  
 ACT GAT GTA GCC TGG TAT CAA CAG AAA CCA  
 10 CGA CAA TCT CCT AAA CTA CTG ATT TAC TCG  
 GCA TCC TAT CGG TAC ACT GGA GTC CCT GAT  
 CGC TTC ACT GGC AGT GGA TCT GGG ACG GAT  
 TTC ACT TTC ACC ATC AGC AGT GTG CAG GCT  
 15 GAA GAC CTG GCA GTT TAT TAC TGT CAG CAA  
 CAT TAT AGT ACT GCG TGG ACG TTC GGT GGT  
 GGC ACC AAG CTG GAA ATC AAA .

23. An Fv region fragment comprising the immunoglobulin H chain variable region fragment according to claim 2 and the immunoglobulin L chain variable region fragment according to claim 12.

24. An Fv region fragment comprising the immunoglobulin H chain variable region fragment according to claim 2 and the immunoglobulin L chain variable region fragment according to claim 13.

25. An Fv region fragment comprising the immunoglobulin H chain variable region fragment according to claim 3 and the immunoglobulin L chain variable region fragment according to claim 14.

26. An Fv region fragment comprising the immunoglobulin H chain variable region fragment according to claim 4 and the immunoglobulin L chain variable region fragment according to claim 15.

27. An Fv region fragment comprising the immunoglobulin H chain variable region fragment according to claim 5 and the immunoglobulin L chain variable region fragment according to claim 16.

FIG. 1

Idio 3

Idio 17

Idio 20

Idio 27

Idio 33

+	$\lambda$	K	G3	G2b	G2a	G1	M	A
+	$\lambda$	K	G3	G2b	G2a	G1	M	A
+	$\lambda$	K	G3	G2b	G2a	G1	M	A
+	$\lambda$	K	G3	G2b	G2a	G1	M	A
+	$\lambda$	K	G3	G2b	G2a	G1	M	A

FIG. 2




FIG. 3




FIG. 4

3 17 20 27 33

FR 1	1	Glu	Glu	Glu	Glu	Glu
	2	Val	Val	Val	Val	Val
	3	Gln	Gln	Lys	Lys	Gln
	4	Leu	Leu	Leu	Leu	Leu
	5	Gln	Gln	Val	Val	Gln
	6	Gln	Gln	Glu	Glu	Gln
	7	Ser	Ser	Ser	Ser	Ser
	8	Gly	Gly	Gly	Gly	Gly
	9	Thr	Thr	Gly	Gly	Ala
	10	Val	Val	Gly	Gly	Glu
	11	Leu	Leu	Leu	Leu	Leu
	12	Ala	Ala	Val	Val	Ala
	13	Arg	Arg	Gln	Gln	Arg
	14	Pro	Pro	Pro	Pro	Pro
	15	Gly	Gly	Gly	Gly	Gly
	16	Ala	Ala	Gly	Gly	Ala
	17	Ser	Ser	Ser	Ser	Ser
	18	Val	Val	Leu	Leu	Val
	19	Lys	Lys	Arg	Arg	Asn
	20	Met	Met	Leu	Leu	Leu
	21	Ser	Ser	Ser	Ser	Ser
	22	Cys	Cys	Cys	Cys	Cys
	23	Lys	Lys	Ala	Ala	Lys
	24	Ala	Ala	Thr	Thr	Ala
	25	Ser	Ser	Ser	Ser	Ser
	26	Gly	Gly	Gly	Gly	Gly
	27	Tyr	Tyr	Leu	Phe	Tyr
	28	Thr	Thr	Thr	Thr	Thr
	29	Phe	Phe	Phe	Phe	Phe
	30	Asn	Asn	Thr	Thr	Thr
CDR 1	31	Ser	Ser	Asp	Asp	Asn
	32	Tyr	Tyr	Tyr	Tyr	Tyr
	33	Trp	Trp	Tyr	Tyr	Trp
	34	Met	Met	Met	Met	Met
FR 2	35	His	His	Asn	Asn	Gln
	36	Trp	Trp	Trp	Trp	Trp
	37	Val	Val	Val	Val	Val
	38	Lys	Lys	Arg	Arg	Lys
	39	Gln	Gln	Gln	Gln	Gln
	40	Arg	Arg	Pro	Pro	Arg
	41	Pro	Pro	Pro	Pro	Pro
	42	Gly	Gly	Gly	Gly	Gly
	43	Gln	Gln	Lys	Lys	Gln
	44	Gly	Gly	Glu	Ala	Gly
	45	Leu	Leu	Leu	Leu	Leu
	46	Glu	Glu	Glu	Glu	Glu
	47	Trp	Trp	Trp	Trp	Trp
	48	Ile	Ile	Leu	Leu	Ile
CDR 2	49	Gly	Gly	Gly	Gly	Gly
	50	Ala	Ala	Phe	Phe	Ala
	51	Ile	Ile	Ile	Ile	Ile
	52	Tyr	Tyr	Arg	Arg	Tyr
	52A	Pro	Pro	Asn	Asn	Pro
	52B	---	---	Lys	Lys	---
	52C	---	---	Ala	Ala	---
	53	Gly	Gly	Asn	Asn	Gly
	54	Asn	Asn	Leu	Tyr	Asp
	55	Ser	Ser	Tyr	Tyr	Gly
	56	Asp	Asp	Thr	Thr	Asp
	57	Ile	Ile	Thr	Thr	Thr
	58	Ser	Ser	Asp	Glu	Arg
	59	Tyr	Tyr	Tyr	Tyr	Tyr
	60	Ser	Ser	Ser	Ser	Thr
	61	Gln	Gln	Ala	Ala	Gln
	62	Asn	Asn	Ser	Ser	Lys
	63	Phe	Phe	Val	Val	Phe
	64	Lys	Lys	Lys	Lys	Lys
	65	Asp	Asp	Gly	Gly	Gly
	66	Arg	Arg	Arg	Arg	Lys
	67	Ala	Ala	Phe	Phe	Ala
FR 3	68	Lys	Lys	Thr	Thr	Thr
	69	Leu	Leu	Ile	Ile	Leu
	70	Thr	Thr	Ser	Ser	Thr
	71	Ala	Ala	Arg	Arg	Ala
	72	Val	Val	Asp	Asp	Ala
	73	Thr	Thr	Asn	Asn	Lys
	74	Ser	Ser	Pro	Ser	Ser
	75	Thr	Thr	Gln	Gln	Ser
	76	Ser	Ser	Ser	Ser	Ser
	77	Thr	Thr	Ile	Ile	Thr
	78	Ala	Ala	Leu	Leu	Ala
	79	Tyr	Tyr	Tyr	Tyr	Tyr
	80	Met	Met	Leu	Leu	Met
	81	Glu	Glu	Gln	Gln	Gln
	82	Leu	Leu	Met	Met	Leu
	82A	Arg	Arg	Asn	Asn	Ser
	82B	Ser	Ser	Thr	Thr	Ser
	82C	Leu	Leu	Leu	Leu	Leu
	83	Thr	Thr	Thr	Arg	Ala
	84	Asn	Asn	Thr	Ala	Ser
	85	Glu	Glu	Glu	Glu	Glu
	86	Asp	Asp	Asp	Asp	Asp
	87	Ser	Ser	Ser	Ser	Ser
	88	Ala	Ala	Ala	Ala	Ala
	89	Val	Val	Thr	Thr	Val
	90	Tyr	Tyr	Tyr	Tyr	Tyr
	91	Phe	Phe	Tyr	Tyr	Tyr
	92	Cys	Cys	Cys	Cys	Cys
	93	Thr	Thr	Ala	Ala	Ala
	94	Lys	Lys	Arg	Arg	Arg
CDR 3	95	Glu	Glu	Asp	Asp	Ser
	96	Glu	Glu	Arg	Gly	Gly
	97	Tyr	Tyr	Gly	Phe	Tyr
	98	Asp	Asp	Gly	Leu	Tyr
	99	Tyr	Tyr	Arg	Arg	Gly
	100	Asp	Asp	Asp	Asp	Ser
	100A	Thr	Thr	---	---	Phe
	100B	---	---	---	---	Val
	100C	---	---	---	---	Gly
	100D	---	---	---	---	---
FR 4	100E	---	---	---	---	---
	100F	---	---	---	---	---
	100G	---	---	---	---	---
	100H	---	---	---	---	---
	100I	---	---	Trp	Trp	---
	100J	---	---	Tyr	Tyr	---
	100K	Leu	Leu	Phe	Phe	Phe
	101	Asp	Asp	Asp	Asp	Ala
	102	Tyr	Tyr	Val	Val	Tyr
	103	Trp	Trp	Trp	Trp	Trp
	104	Gly	Gly	Gly	Gly	Gly
	105	Gln	Gln	Ala	Ala	Gln
	106	Gly	Gly	Gly	Gly	Gly
	107	Thr	Thr	Thr	Thr	Thr
	108	Ser	Ser	Thr	Thr	Leu
	109	Val	Val	Val	Val	Val
	110	Thr	Thr	Thr	Thr	Thr
	111	Val	Val	Val	Val	Val
	112	Ser	Ser	Ser	Ser	Ser
	113	Ser	Ser	Ser	Ser	Ala

## FIG. 5

3 17 20 27 33

F R 1	1	Asp	Asp	Asp	Asp	Asp
	2	Ile	Ile	Ile	Ile	Ile
	3	Val	Val	Val	Val	Val
	4	Leu	Leu	Leu	Met	Met
	5	Thr	Thr	Thr	Thr	Thr
	6	Gln	Gln	Gln	Gln	Gln
	7	Ser	Ser	Ser	Ser	Ser
	8	Pro	Pro	Pro	His	His
	9	Ala	Ala	Ala	Lys	Lys
	10	Ser	Ser	Ser	Phe	Phe
	11	Leu	Leu	Leu	Met	Met
	12	Ala	Ala	Ala	Ser	Ser
	13	Val	Val	Val	Thr	Thr
	14	Ser	Ser	Ser	Ser	Ser
	15	Pro	Leu	Leu	Val	Val
	16	Leu	Gly	Gly	Gly	Gly
	17	Gly	Gln	Gln	Asp	Asp
	18	Gln	Arg	Arg	Arg	Arg
	19	Arg	Ala	Ala	Val	Val
	20	Ala	Ser	Thr	Ser	Thr
	21	Thr	Ile	Ile	Ile	Ile
	22	Ile	Ser	Ser	Thr	Thr
	23	Ser	---	---	Cys	Cys
C D R 1	24	Tyr	Tyr	Tyr	Lys	Lys
	25	Arg	Arg	Arg	Ala	Ala
	26	Ala	Ala	Ala	Ser	Ser
	27	Ser	Ser	Ser	Gln	Gln
	27A	Lys	Lys	Lys	---	---
	27B	Ser	Ser	Ser	---	---
	27C	Val	Val	Val	---	---
	27D	Gln	Ser	Ser	---	---
	27E	Leu	Thr	Thr	---	---
	27F	His	---	---	---	---
	28	Leu	Ser	Ser	Asp	Asp
	29	Ala	Gly	Gly	Val	Val
F R 2	30	Ile	Tyr	Tyr	Asn	Thr
	31	Val	Ser	Ser	Thr	Thr
	32	Tyr	Tyr	Tyr	Ala	Asp
	33	Met	Met	Met	Val	Val
	34	His	His	His	Ala	Ala
	35	Trp	Trp	Trp	Trp	Trp
	36	Asn	Asn	Asn	Tyr	Tyr
	37	Gln	Gln	Gln	Gln	Gln
	38	Gln	Gln	Gln	Gln	Gln
	39	Lys	Lys	Arg	Lys	Lys
	40	Pro	Pro	Pro	Pro	Pro
	41	Gly	Gly	Gly	Gly	Arg
C D R 2	42	Gln	Gln	Gln	Gln	Gln
	43	Pro	Pro	Pro	Ser	Ser
	44	Pro	Pro	Pro	Pro	Pro
	45	Arg	Arg	Arg	Lys	Lys
	46	Leu	Leu	Leu	Leu	Leu
	47	Leu	Leu	Leu	Leu	Leu
	48	Ile	Ile	Ile	Leu	Ile
	49	Tyr	Tyr	Tyr	Tyr	Tyr
	50	Leu	Leu	Leu	Ser	Ser
	51	Val	Val	Val	Ala	Ala
	52	Ser	Ser	Ser	Ser	Ser
	53	Asn	Asn	Asn	Tyr	Tyr
	54	Leu	Leu	Leu	Arg	Arg
	55	Glu	Glu	Asp	Tyr	Tyr
	56	Ser	Ser	Ser	Thr	Thr
	57	Gly	Gly	Gly	Gly	Gly
	58	Val	Val	Val	Val	Val
	59	Pro	Pro	Pro	Pro	Pro
	60	Ala	Ala	Ala	Asp	Asp
	61	Arg	Arg	Arg	His	Arg
	62	Phe	Phe	Phe	Phe	Phe
	63	Ser	Ser	Ser	Thr	Thr
	64	Gly	Gly	Gly	Gly	Gly

F R 3	65	Ser	Ser	Ser	Ser	Ser
	66	Gly	Gly	Gly	Gly	Gly
	67	Gly	Ser	Ser	Ser	Ser
	68	Gly	Gly	Gly	Gly	Gly
	69	Thr	Thr	Thr	Thr	Thr
	70	Asp	Asp	Asp	Asp	Asp
	71	Phe	Phe	Phe	Phe	Phe
	72	Thr	Thr	Thr	Thr	Thr
	73	Leu	Leu	Leu	Phe	Phe
	74	Asn	Asn	Asn	Thr	Thr
	75	Ile	Ile	Ile	Ile	Ile
	76	His	His	His	Ser	Ser
	77	Pro	Pro	Pro	Ser	Ser
	78	Val	Val	Val	Val	Val
	79	Glu	Glu	Glu	Gln	Gln
	80	Glu	Glu	Glu	Ala	Ala
	81	Glu	Glu	Glu	Glu	Glu
	82	Asp	Asp	Asp	Asp	Asp
	83	Ala	Ala	Ala	Leu	Leu
	84	Ala	Ala	Ala	Ala	Ala
	85	Thr	Thr	Thr	Val	Val
C D R 3	86	Tyr	Tyr	Tyr	Tyr	Tyr
	87	Tyr	Tyr	Tyr	Tyr	Tyr
	88	Cys	Cys	Cys	Cys	Cys
	89	Gln	Gln	Gln	Gln	Gln
	90	His	His	His	Gln	Gln
	91	Ile	Ile	Ile	His	His
	92	Arg	Arg	Glu	Tyr	Tyr
	93	Val	Gly	Gly	Ser	Ser
	94	Ala	Ala	Ala	Pro	Thr
	95	---	---	---	Pro	Ala
	95A	---	---	---	---	---
	95B	---	---	---	---	---
	95C	---	---	---	---	---
	95D	---	---	---	---	---
	95E	---	---	---	---	---
	95F	---	---	---	---	---
F R 4	96	Tyr	Tyr	Tyr	Leu	Trp
	97	Thr	Thr	Thr	Thr	Thr
	98	Phe	Phe	Phe	Phe	Phe
	99	Gly	Gly	Gly	Gly	Gly
	100	Gly	Gly	Gly	Ala	Gly
	101	Gly	Gly	Gly	Gly	Gly
	102	Thr	Thr	Thr	Thr	Thr
	103	Lys	Lys	Lys	Lys	Lys
	104	Leu	Leu	Leu	Leu	Leu
	105	Glu	Glu	Glu	Glu	Glu
	106	Ile	Ile	Ile	Leu	Ile
	106A	---	---	---	---	---
	107	Lys	Lys	Lys	Lys	Lys



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number  
EP 94 11 5683

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.4)
A	US-A-5 208 146 (R. ERIE) * the whole document *	1-27	C12N15/13 C07K16/42
A	WO-A-89 00050 (AKZO NV) * claims * * examples *	1-27	
A	WO-A-93 10221 (THE REGENTS OF THE UNIVERSITY OF CALIFORNIA) * the whole document *	1-27	
A	EUROPEAN JOURNAL OF CANCER AND CLINICAL ONCOLOGY, vol.24, no.5, May 1988, OXFORD, GB pages 829 - 838 Y. AOTSUKA ET AL. 'Identification of a malignant cell associated antigen recognized by a human monoclonal antibody.' * abstract *	1-27	
A	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE USA, vol.80, no.20, October 1983, WASHINGTON DC, USA pages 6327 - 6331 M. GLASSY ET AL. 'UC 729-6, a human lymphoblastoid B-cell line useful for generating antibody-secreting human-human hybridomas.' * abstract *	1-27	TECHNICAL FIELDS SEARCHED (Int.Cl.4) C12N C07K
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 16 March 1995	Examiner Nooij, F
<p><b>CATEGORY OF CITED DOCUMENTS</b></p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons A : member of the same patent family, corresponding document</p>			

EPO FORM 150 (03.03) (P04C01)





European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number  
EP 94 11 5683

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
A	CANCER RESEARCH, vol.52, no.9, 1 May 1992, PHILADELPHIA PA, USA pages 2603 - 2609 W. TADDEI-PETERS ET AL. 'Quantitation of human tumor-reactive monoclonal antibody 16.88 in the circulation and localization of 16.88 in colorectal metastatic tumor tissue using murine antiidiotypic antibodies.' * abstract *	1-27	
P,A	MOLECULAR IMMUNOLOGY, vol.30, no.16, November 1993, OXFORD, GB pages 1481 - 1489 K. YAGO ET AL. 'Immunoglobulin variable region sequences of two human monoclonal antibodies directed to an onco-developmental carbohydrate antigen, lactotetraosylceramide (LcOse4Cer).' * abstract *	1-27	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
The present search report has been drawn up for all claims.			
Place of search THE HAGUE		Date of completion of the search 16 March 1995	Examiner Nooij, F.
CATEGORY OF CITED DOCUMENTS		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons A : technological background P : non-written disclosure I : intermediate document	
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background P : non-written disclosure I : intermediate document		A : member of the same patent family, corresponding document	

EPO FORM 1503 (3.93) (P/M/C/N)

[Faint, mostly illegible text covering the majority of the page, appearing to be a multi-paragraph document.]

OFFICE/JCWS

MAY - 3 2012

RECEIVED